

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 211/90, 409/04, 407/04, 401/04, A61K 31/44		A2	(11) International Publication Number: WO 97/27177
			(43) International Publication Date: 31 July 1997 (31.07.97)
(21) International Application Number: PCT/US97/01252		(US). JIANG, Ji-Long [CN/US]; 10802 Drawbridge Court, Silver Spring, MD 20902 (US). KIM, Yong-Chul [KR/US]; Apartment 406, 236 Congressional Lane, Rockville, MD 20852 (US). KARTON, Yishai [IL/IL]; 27/16 Bailik Street, Nes Ziona (IL). VAN RHEE, Albert, M. [NL/US]; Apartment 1343, 10900 Bucknell Drive, Silver Spring, MD 20902 (US). (74) Agents: KILYK, John, Jr. et al.; Leydig, Voit & Mayer, Ltd., Two Prudential Plaza, Suite 4900, 180 North Stetson, Chicago, IL 60601-6780 (US). (81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>	
(22) International Filing Date: 29 January 1997 (29.01.97)			
(30) Priority Data: 60/010,737 29 January 1996 (29.01.96) US 60/021,191 3 July 1996 (03.07.96) US			
(60) Parent Applications or Grants (63) Related by Continuation US 60/021,191 (CIP) Filed on 3 July 1996 (03.07.96) US 60/010,737 (CIP) Filed on 29 January 1996 (29.01.96)			
(71) Applicant (for all designated States except US): THE UNITED STATES OF AMERICA, represented by THE SECRETARY, DEPARTMENT OF HEALTH AND HUMAN SERVICES National Institutes of Health, Office of Technology Transfer, Box OTT [US/US]; Bethesda, MO 20892-9902 (US).			
(72) Inventors; and (75) Inventors/Applicants (for US only): JACOBSON, Kenneth, A. [US/US]; 11606 Fulham Street, Silver Spring, MD 20902			
(54) Title: DIHYDROPYRIDINE-, PYRIDINE-, BENZOPYRAN- ONE- AND TRIAZOLOQUINAZOLINE DERIVATIVE, THEIR PREPARATION AND THEIR USE AS ADENOSINE RECEPTOR ANTAGONISTS			
(57) Abstract			
<p>The present invention provides certain novel compounds, compositions, and a method of treating a mammal by blocking its adenosine receptors comprising administering at least one compound of the present invention. Examples of the present inventive compounds include certain flavonoids of formulae (I) and (II), wherein R₁ to R₄ are as defined in the description, and M is -CH(OH)-CH(R₂) or -C(OH)=C(R₂)- and R₁, R₂ are as defined in the description; or dihydropyridines of formula (III), wherein R₂ to R₆ are as defined in the description; or pyridines of formula (IV), wherein R₂ to R₆ are as defined in the description, or triazoloquinazolines of formula (V) wherein R₁ and R₂ are as defined in the description; and their derivatives, or pharmaceutically acceptable salts thereof.</p>			
<div style="display: flex; justify-content: space-around; align-items: flex-end;"> <div style="text-align: center;"> (I) </div> <div style="text-align: center;"> (II) </div> <div style="text-align: center;"> (III) </div> <div style="text-align: center;"> (IV) </div> <div style="text-align: center;"> (V) </div> </div>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LJ	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

DIHYDROPYRIDINE-, PYRIDINE-, BENZOPYRAN- ONE- AND TRIAZOLOQUINAZOLINE
DERIVATIVE, THEIR PREPARATION AND THEIR USE AS ADENOSINE RECEPTOR
ANTAGONISTS

CROSS-REFERENCE TO RELATED APPLICATIONS

This is a continuation-in-part application of United
States provisional patent application Serial No.
60/010,737, filed January 29, 1996, and Serial No. 60/021,
191, filed July 3, 1996, the disclosures of which are
incorporated herein in their entirety by reference.

TECHNICAL FIELD OF THE INVENTION

The present invention relates to adenosine receptor
antagonists, pharmaceutical compositions, and methods of
selectively blocking adenosine receptors in a mammal. The
present invention also relates to methods of treating
various medical disorders with adenosine receptor
antagonists.

BACKGROUND OF THE INVENTION

The use of caffeine and other alkylxanthines as
physiological stimulants is well known. The principle
mechanism by which caffeine and other alkylxanthines act
as physiological stimulants is by blocking the effects of
the ubiquitous neuromodulator adenosine. Daly, "Mechanism
of action of caffeine", in Caffeine, Coffee and Health,
(S. Garattini, ed), Chapter 4, pp. 97-150 (1993).
Adenosine is produced locally in response to increased
activity or stress to the system. This feedback mechanism
allows the organ to compensate for the stress by
decreasing energy demand (depressant activity) and
increasing oxygen supply (e.g., by vasodilation). Bruns,
Nucleosides & Nucleotides, 10, 931-944 (1991).

Adenosine plays several key physiological roles. In
addition to its role in intermediary metabolism, adenosine
displays a number of receptor-mediated physiological
actions, including dilation of coronary vessels,
inhibition of platelet aggregation, and inhibition of
lipolysis. Bruns et al., Proc. Nat. Acad. Sci. U.S.A., 77,

5547-5551 (1980). Adenosine receptors, belonging to the superfamily of the G protein-coupled receptors, are generally divided into two major subclasses, A_1 and A_2 , on the basis of the differential affinities of a number of adenosine receptor agonists and antagonists for the receptors, their primary structures, and the secondary messenger systems to which they couple. Thus, A_2 receptors, which can be further subdivided into A_{2a} and A_{2b} , stimulate adenylate cyclase, whereas A_1 receptors may couple to a variety of secondary messenger systems, including those involved in the inhibition of adenylate cyclase, the inhibition or stimulation of phosphoinositol turnover, the activation of guanylate cyclase, the activation of potassium influx, and the inhibition of calcium influx (van Galen et al., Med. Res. Rev., **12**, 423-471 (1992); Jacobson et al., J. Med. Chem., **35**, 407-422 (1992)).

Recently, a novel adenosine receptor was identified on the basis of its primary structure and cloned from rat brain (Zhou et al., Proc. Natl. Acad. Sci. U.S.A., **89**, 7432-7436 (1992)) and rat testes (Meyerhof et al., FEBS Lett., **284**, 155-160 (1991)). The putative transmembrane domains of the novel adenosine receptor, which has been designated the A_3 receptor, show 58% identity with the canine A_1 receptor and 57% with the canine A_{2a} receptor. Like the A_1 receptor, the A_3 receptor is negatively coupled to adenylate cyclase (Zhou et al.).

The distribution of the A_3 receptor is found primarily in the central nervous system (CNS) (Zhou et al.), brain, testes (Meyerhof et al.), and immune system, where it appears to be involved in the modulation of release from mast cells of mediators of the immediate hypersensitivity reaction (Ramkumar et al., J. Biol. Chem., **268**, 16887-16890 (1993)). It is believed that A_3 -selective compounds will have utility in the therapeutic and/or prophylactic treatment of cardiac disease, infertility, kidney disease, and CNS disorders.

It is further believed that selective A₃-adenosine receptor antagonists should serve as cerebroprotective, anti-asthmatic, or anti-inflammatory agents. Beaven et al., Trends Pharmacol. Sci., **15**, 13-4 (1994); Jacobson et al., Drugs of the Future, **20**, 689-699 (1995); von Lubitz et al., Eur. J. Pharmacol., **275**, 23-29 (1995).


Copending U.S. patent applications Serial No. 08/274,628, filed July 13, 1994, and Serial No. 08/396,111, filed February 28, 1995, disclose certain A₃ selective agonists, particularly N⁶-benzyladenosine-5'-uronamide and related substituted compounds, xanthine riboside derivatives, pharmaceutical compositions comprising such compounds, and the method of use of such compounds.

The foregoing indicates that there is a need for antagonists for adenosine receptors. The present invention seeks to provide such compounds, as well as methods of using these compounds to selectively block adenosine receptors in mammals, and pharmaceutical compositions comprising such compounds. These and other objects and advantages of the present invention, as well as additional inventive features, will be apparent from the description of the invention provided herein.

25

SUMMARY OF THE INVENTION

The present invention provides a compound of the formula I




(I)

30

wherein R₁ and R₂ are selected from the group consisting of hydrogen, hydroxy, C₁-C₆ alkyloxy, and C₁-C₆ alkylcarbonyloxy; R₃ is selected from the group consisting

of hydrogen, hydroxy, C_1 - C_6 alkyloxy, and C_2 - C_6 alkenyloxy, said alkenyloxy together with the carbon atom of the phenyl ring forming an oxygen heterocycle; and R_4 is selected from the group consisting of phenyl, styryl, phenylbutadienyl, phenylacetylenyl, and -CH=N-phenyl, and substituted phenyl, styryl, phenylacetylenyl, and phenylbutadienyl, wherein the phenyl ring is substituted with 1 to 5 C_1 - C_6 alkyloxy groups; with the provisos that when R_3 is hydrogen, R_1 and R_2 are neither hydroxy nor alkyloxy; when R_1 , R_2 , and R_3 are hydrogen, R_4 is neither phenyl nor alkyloxyphenyl; when R_3 is hydrogen and R_4 is phenyl, neither R_1 nor R_2 is alkylcarbonyloxy; and when R_3 is hydroxy or alkyloxy, R_1 and R_2 are not dihydroxy.


The present invention further provides a compound of the formula II



(II)

wherein R_1 is selected from the group consisting of hydroxyl and C_1 - C_6 alkoxy, and M is a divalent radical selected from the group consisting of -CH(OH)-CH(R_2)- and -C(OH)=C(R_2)-, wherein R_2 is selected from the group consisting of styryl and phenylacetylenyl.

The present invention provides a compound of the formula III




(III)

or a pharmaceutically acceptable salt thereof, wherein R_2 is a C_1 - C_6 alkyl; R_6 is selected from the group consisting of C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, and phenyl which may be further substituted with C_1 - C_6 alkyl, halo, nitro, furyl,


or thienyl; R_1 is selected from the group consisting of C_1 - C_6 alkyl, C_1 - C_6 alkyloxycarbonyl, aryl C_1 - C_6 alkyloxycarbonyl, C_1 - C_6 alkylthiocarbonyl, C_1 - C_6 alkylaminocarbonyl, and C_1 - C_6 alkyloxy C_1 - C_6 alkylcarbonyl, or R_1 together with R_2 forms a ring having 2-4 methylene groups, and C_1 - C_6 alkenyloxycarbonyl; R_4 is selected from the group consisting of C_1 - C_6 alkyl, aryl C_2 - C_6 alkenyl, C_1 - C_6 alkylamino, C_1 - C_6 alkyl silyl C_1 - C_6 alkyloxy, aryl, heterocyclic, aryl C_1 - C_6 alkyl, phenylacetylenyl which may be further substituted with nitro, C_1 - C_6 alkyl, hydroxy, halo, amino, carboxy, C_1 - C_6 alkoxy, C_1 - C_6 haloalkyl, or C_1 - C_6 alkylamino, and styryl whose phenyl ring may be further substituted with one or more substituents selected from the group consisting of halo, nitro, amino, hydroxy, C_1 - C_6 alkyl, cyano, C_1 - C_6 alkyloxy, C_1 - C_6 alkyloxycarbonyl, C_1 - C_6 alkylcarbonyl, hydroxy C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, carboxy, aminocarbonyl, C_1 - C_6 alkylamino, amino C_1 - C_6 alkyl, and C_1 - C_6 dialkylamino; and R_5 is selected from the group consisting of C_1 - C_6 alkyloxycarbonyl, aryl C_1 - C_6 alkyloxycarbonyl, C_1 - C_6 alkyloxy C_1 - C_6 alkyloxycarbonyl, aryloxy C_1 - C_6 alkyloxycarbonyl, C_1 - C_6 alkyloxycarbonyl, aryl C_1 - C_6 alkyloxy C_1 - C_6 alkyloxycarbonyl, silyl C_1 - C_6 alkyloxycarbonyl, C_1 - C_6 alkylthio, hydroxy, and C_1 - C_6 alkylamino, wherein said aryl may be further substituted with C_1 - C_6 alkyl, C_1 - C_6 halo alkyl, trifluoromethyl, halo, nitro, C_1 - C_6 amino alkyl, C_1 - C_6 aminoalkylamino, or C_1 - C_6 amino alkylamino carbonyl; wherein said aryl is phenyl or naphthyl.

The present invention further provides a compound of the formula IV



or pharmaceutically acceptable salts thereof, wherein R_2 is selected from the group consisting of hydrogen and C_1 - C_6 alkyl; R_3 is selected from the group consisting of hydrogen

- and C₁-C₆ alkyloxycarbonyl; R₄ is selected from the group consisting of C₁-C₆ alkyl, phenyl C₂-C₆ alkenyl, phenyl C₂-C₆ alkynyl, aryl, and aryl substituted with one or more substituents selected from the group consisting of nitro and C₁-C₆ alkyloxy; R₅ is selected from the group consisting of hydrogen, C₁-C₆ alkyloxycarbonyl, and aryl C₁-C₆ alkyloxy carbonyl; R₆ is selected from the group consisting of hydrogen, aryl, and C₁-C₆ alkyl; with the proviso that when R₂=R₃=R₅=R₆=hydrogen, R₄ is not alkyl.
- The present invention further provides a compound of the formula V



(V)

- wherein R₁ is selected from the group consisting of C₁-C₆ alkylcarbonyl, C₁-C₆ alkyloxycarbonyl, amino C₁-C₆ alkylcarbonyl, and arylcarbonyl wherein the aryl may be further substituted with halo, nitro, hydroxy, amino or cyano; and R₂ is hydrogen or halogen.

- The present invention further provides pharmaceutical compositions comprising any of the aforesaid compounds and a method of treating a mammal comprising selectively blocking one or more of the adenosine receptors of the mammal by administering to the mammal at least one compound of formulas I-V.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 depicts a method of synthesis of alkylated derivatives of galangin and morin. Alkyl iodide was used as the alkylating agent for the preparation of 2, 3a, 3b, 4, and 6. Ethyl bromide was used as the alkylating agent for the preparation of 7, and alkyl sulfate was used as the alkylating agent for the preparation of 8.

Figure 2 depicts a method of synthesis of the 2-phenylacetylenyl and 2-cinnamyl derivatives, 10, 15, and 16, starting from 4-methoxyfuranochromone.

Figure 3 depicts a method of synthesis of certain 2-substituted flavonoid derivatives, 19-24, starting from visnagin.

Figure 4 depicts the structures of flavonoid derivatives 41-44.

Figure 5 depicts the structures of flavonoid derivatives 45-51.

Figure 6 depicts a method of synthesis of certain 1,4-dihydropyridine derivatives.

Figure 7 depicts a method of oxidation of certain 1,4-dihydropyridine derivatives.

Figure 8 depicts a representative set of competition curves for inhibition binding of [¹²⁵I]AB-MECA by compounds 12 (triangles), 14 (circles), and 18 (diamonds), at human brain A₁ receptors expressed in human HEK-293 cells at 25°C.

Figure 9A depicts a part of the reaction scheme for the synthesis of a diastereomeric pair of a 5-ester substituted 1,4-dihydropyridine.

Figure 9B depicts another part of the reaction scheme for the synthesis of a diastereomeric pair of a 5-ester substituted 1,4-dihydropyridine.

Figure 10A depicts the percentage of HL-60 apoptotic cells (vertical axis) as a function of the concentration (horizontal axis) of A₁ adenosine receptor agonists IB-MECA

(°) and Cl-IB-MECA (•), as determined by fluorescent cell sorting (flow cytometric DNA analysis).

Figure 10B depicts the percentage of U-937 apoptotic cells (vertical axis) as a function of the concentration (horizontal axis) of A₁ adenosine receptor agonists IB-MECA (°) and Cl-IB-MECA (•), as determined by fluorescent cell sorting (flow cytometric DNA analysis).

Figure 11A depicts the number of living HL-60 cells (vertical axis) as a function of the concentration (horizontal axis) of A₁ adenosine receptor antagonist and a low concentration of Cl-IB-MECA. The antagonist was compound 101. In Figures 11A-D, for each set of curves, control (--), antagonist alone (Δ), or antagonist in the presence of 10nM (°) or 1 μM (•) Cl-IB-MECA are shown.

Figure 11B depicts the number of living HL-60 cells (vertical axis) as a function of the concentration (horizontal axis) of the A₁ adenosine receptor antagonist and low concentration of Cl-IB-MECA. The antagonist was compound L-249313(6-carboxymethyl-5,9-dihydro-9-methyl-2-phenyl-[1,2,4]triazolo[5,1-a][2,7]-naphthyridine).

Figure 11C depicts the number of living U-937 cells (vertical axis) as a function of the concentration (horizontal axis) of the A₁ adenosine receptor antagonist and a low concentration of Cl-IB-MECA. The antagonist was compound 101.

Figure 11D depicts the number of living U-937 cells (vertical axis) as a function of the concentration (horizontal axis) of the A₁ adenosine receptor antagonist and a low concentration of Cl-IB-MECA. The antagonist was compound L-249313.


Figure 12 depicts the percent change in dynamic compliance (C_{dyn} , vertical axis) of rabbit lung as a function of the adenosine agonist concentration using Cl-IB-MECA (o), APNEA (), or IB-MECA (Δ). (•) represents percent change in C_{dyn} as a result of exposure to compound 101 plus Cl-IB-MECA and (\blacktriangle) represents percent change in C_{dyn} as a result of exposure to compound 101 plus IB-MECA.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

The present invention may be best understood with reference to the accompanying drawings and to the following detailed description of the preferred embodiments.

The present invention provides certain derivatives of flavonoids, dihydropyridines, pyridines, and triazoloquinazolines, suitable for blocking one or more of the adenosine receptors of a mammal such as human, as set forth herebelow in greater detail.

The present invention provides a compound of the formula I



wherein R_1 and R_3 are selected from the group consisting of hydrogen, hydroxy, C_1 - C_6 alkyloxy, and C_1 - C_6 alkylcarbonyloxy; R_2 is selected from the group consisting of hydrogen, hydroxy, C_1 - C_6 alkyloxy, and C_2 - C_6 alkenyloxy, said alkenyloxy together with the carbon atom of the phenyl ring forming an oxygen heterocycle; and R_4 is selected from the group consisting of phenyl, styryl, phenylbutadienyl, phenylacetylenyl, and $-CH=N$ -phenyl, and substituted phenyl, styryl, phenylacetylenyl, and phenylbutadienyl, wherein the phenyl ring is substituted with 1 to 5 C_1 - C_6 alkyloxy groups; with the provisos that

when R_3 is hydrogen, R_1 and R_2 are neither hydroxy nor alkoxy; when R_1 , R_2 , and R_3 are hydrogen, R_4 is neither phenyl nor alkyloxyphenyl; when R_3 is hydrogen and R_4 is phenyl, neither R_1 nor R_2 is alkylcarbonyloxy; and when R_3 is hydroxy or alkoxy, R_1 and R_2 are not dihydroxy.

Particular embodiments include a compound of formula I wherein (a) R_4 is phenyl, (b) R_4 is phenyl and R_3 is a C_1 - C_3 alkoxy, (c) R_4 is phenyl, R_3 is a C_1 - C_3 alkoxy, and R_1 and R_2 are 5,7-di(C_1 - C_3 alkoxy), (d) R_4 is a 2,4-di(C_1 - C_3 alkoxy)phenyl, (e) R_3 is a C_1 - C_3 alkoxy and R_4 is a 2,4-di(C_1 - C_3 alkoxy)phenyl, (f) R_3 is a C_1 - C_3 alkoxy, R_4 is a 2,4-di(C_1 - C_3 alkoxy)phenyl, and R_1 and R_2 are the same and are selected from the group consisting of methoxy and ethoxy, (g) R_1 is 5-hydroxy and R_2 is one of methoxy and ethoxy, (h) R_4 is phenylacetylenyl, (i) R_4 is phenylacetylenyl and R_3 is hydroxy, and (j) R_4 is phenylacetylenyl, R_3 is hydroxy, and one of R_1 and R_2 is methoxy.

The present invention further provides a compound of formula I, wherein the compound is a 4-(C_1 - C_6 alkoxy)-7-styrylvisnagin. Examples of such compounds include 4-methoxy-7-trans-styrylvisnagin, 4-ethoxy-7-trans-styrylvisnagin, and 4-propoxy-7-trans-styrylvisnagin.


Other particular embodiments include compounds of formula I, wherein the compound is a C_1 - C_6 alkoxy-7-phenylbutadienylvisnagin. Examples of such compounds include 4-methoxy-7-phenylbutadienylvisnagin and 4-ethoxy-7-phenylbutadienylvisnagin.

The present invention further provides a compound of formula I, wherein the compound is a C_1 - C_6 alkoxy-7-(CH=N-phenyl)visnagin. An example of such compound is 4-methoxy-7-(CH=N-phenyl)visnagin.

The present invention further provides a compound of formula I, wherein the compound is selected from the group consisting of 3,5,7-triacetoxyflavone, 3,5,7-trimethoxyflavone, 3,5,7-triethoxyflavone, 3,7-diethoxy-5-hydroxyflavone, 3,5,7-tripropoxyflavone, 3,4',5,7-

tetramethoxyflavone, 2',3,4',7-tetraethoxy-5-hydroxyflavone, 2',3,4',5,7-pentamethoxyflavone, 2',3,4',5,7-pentaethoxyflavone, hexamethylmyricetin, and 3-hydroxy-4'-phenylacetylenyl-6-methoxyflavone.

- 5 The present invention further provides a compound of the formula II



- wherein R_1 is selected from the group consisting of hydroxyl and C_1 - C_6 alkoxy, and M is a divalent radical
 10 selected from the group consisting of $-CH(OH)-CH(R_2)-$ and $-C(OH)=C(R_2)-$, wherein R_2 is selected from the group consisting of styryl and phenylacetylenyl.

- Examples of compounds of formula II include 2-phenylacetylenyl-3-hydroxy-6-methoxyflavone, *trans*-2-styryl-3-hydroxy-6-methoxyflavone, and *trans*-2-phenylacetylenyl-3-hydroxy-6-methoxyflavone.
 15

The present invention further provides a compound of the formula III




- 20 or a pharmaceutically acceptable salt thereof, wherein R_2 is a C_1 - C_6 alkyl; R_5 is selected from the group consisting of C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, and phenyl which may be further substituted with C_1 - C_6 alkyl, halo, nitro, furyl, or thienyl; R_3 is selected from the group consisting of C_1 - C_6 alkyl, C_1 - C_6 alkyloxycarbonyl, aryl C_1 - C_6 alkyloxycarbonyl, C_1 - C_6 alkylthiocarbonyl, C_1 - C_6 alkylaminocarbonyl, and C_1 - C_6 alkyloxy C_1 - C_6 alkylcarbonyl, or R_3 together with R_2 forms a ring having 2-4 methylene
 25 groups, and C_1 - C_6 alkenyloxycarbonyl; R_4 is selected from the group consisting of C_1 - C_6 alkyl, aryl C_2 - C_6 alkenyl, C_1 -
 30

- C₆ alkylamino, C₁-C₆ alkyl silyl C₁-C₆ alkyloxy, aryl, heterocyclic, aryl C₁-C₆ alkyl, phenylacetylenyl which may be further substituted with nitro, C₁-C₆ alkyl, hydroxy, halo, amino, carboxy, C₁-C₆ alkoxy, C₁-C₆ haloalkyl, or C₁-C₆ alkylamino, and styryl whose phenyl ring may be further substituted with one or more substituents selected from the group consisting of halo, nitro, amino, hydroxy, C₁-C₆ alkyl, cyano, C₁-C₆ alkyloxy, C₁-C₆ alkyloxycarbonyl, C₁-C₆ alkylcarbonyl, hydroxy C₁-C₆ alkyl, C₁-C₆ haloalkyl, carboxy, aminocarbonyl, C₁-C₆ alkylamino, amino C₁-C₆ alkyl, and C₁-C₆ dialkylamino; and R₅ is selected from the group consisting of C₁-C₆ alkyloxycarbonyl, aryl C₁-C₆ alkyloxycarbonyl, C₁-C₆ alkyloxy C₁-C₆ alkyloxycarbonyl, aryloxy C₁-C₆ alkyloxycarbonyl, C₁-C₆ alkyloxycarbonyl, aryl C₁-C₆ alkyloxycarbonyl, C₁-C₆ alkyloxycarbonyl, silyl C₁-C₆ alkyloxycarbonyl, C₁-C₆ alkylthio, hydroxy, and C₁-C₆ alkylamino, wherein the aryl moiety of said R₅ may be further substituted with C₁-C₆ alkyl, C₁-C₆ halo alkyl, trifluoromethyl, halo, nitro, C₁-C₆ amino alkyl, C₁-C₆ aminoalkylamino, or C₁-C₆ amino alkylamino carbonyl; wherein the aryl moiety of said R₃, R₄, R₅, and R₆ is independently phenyl or naphthyl.

- Particular embodiments include compounds of formula III, wherein (a) R₂ is methyl, (b) R₂ is methyl and R₃ is selected from the group consisting of methoxycarbonyl and ethoxycarbonyl, (c) R₂ is methyl, R₃ is selected from the group consisting of methoxycarbonyl and ethoxycarbonyl, and R₆ is selected from the group consisting of C₁-C₄ alkyl and phenyl, (d) R₂ is methyl, R₃ is selected from the group consisting of methoxycarbonyl and ethoxycarbonyl, R₆ is selected from the group consisting of C₁-C₄ alkyl and phenyl, and R₄ is selected from the group consisting of C₁-C₃ alkyl, (e) R₂ is methyl, R₃ is selected from the group consisting of methoxycarbonyl and ethoxycarbonyl, R₆ is selected from the group consisting of C₁-C₄ alkyl and phenyl, R₄ is selected from the group consisting of C₁-C₃ alkyl, and R₅ is selected from the group consisting of

- methoxycarbonyl, ethoxycarbonyl, methoxyethoxycarbonyl, and benzyloxycarbonyl, (f) R_2 is methyl, R_3 is selected from the group consisting of methoxycarbonyl and ethoxycarbonyl, and R_4 is phenyl substituted with one or more substituents selected from the group consisting of nitro, trifluoromethyl, methoxy, hydroxy, and methylenedioxy, (g) R_2 is methyl, R_3 is selected from the group consisting of methoxycarbonyl and ethoxycarbonyl, R_4 is phenyl substituted with one or more substituents selected from the group consisting of nitro, trifluoromethyl, methoxy, hydroxy, and methylenedioxy, and R_5 is selected from the group consisting of methoxycarbonyl, ethoxycarbonyl, and methoxyethoxycarbonyl, (h) R_2 is methyl, R_3 is ethoxycarbonyl, R_4 is phenylacetylenyl, R_5 is benzyloxycarbonyl which may be further substituted with methyl, trifluoromethyl, halo, iodo, or nitro groups, and R_6 is phenyl, (i) R_2 is methyl, R_3 and R_5 are ethoxycarbonyl, R_4 is 2-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-furyl, or 2-benzofuryl, and R_6 is phenyl, (j) R_2 is methyl, R_3 is ethoxycarbonyl, R_4 is phenylacetylenyl, R_5 is phenylethoxy or phenylpropoxy, and R_6 is phenyl, and (k) R_2 is methyl, R_3 is ethoxycarbonyl, R_4 is phenylacetylenyl, R_5 is thioethoxy, and R_6 is phenyl.
- 25 The present invention further provides a compound of the formula IV




- or pharmaceutically acceptable salts thereof, wherein R_2 is selected from the group consisting of hydrogen and C_1 - C_6 alkyl; R_3 is selected from the group consisting of hydrogen and C_1 - C_6 alkyloxycarbonyl; R_4 is selected from the group consisting of C_1 - C_6 alkyl, phenyl C_2 - C_6 alkenyl, phenyl C_2 - C_6 alkynyl, aryl, and aryl substituted with one or more substituents selected from the group consisting of nitro
- 30

and C₁-C₆ alkyloxy; R₅ is selected from the group consisting of hydrogen, C₁-C₆ alkyloxycarbonyl, and aryl C₁-C₆ alkyloxy carbonyl; and R₆ is selected from the group consisting of hydrogen, aryl, and C₁-C₆ alkyl; with the
5 proviso that when R₂=R₃=R₅=R₆=hydrogen, R₄ is not alkyl.

Particular embodiments include compounds of formula IV, wherein (a) R₂ is selected from the group consisting of hydrogen and methyl, (b) R₂ is selected from the group consisting of hydrogen and methyl, and R₃ and R₅ are same
10 or different and selected from the group consisting of hydrogen, methoxycarbonyl, and ethoxycarbonyl, (c) R₂ is selected from the group consisting of hydrogen and methyl, R₃ and R₅ are same or different and selected from the group consisting of hydrogen, methoxycarbonyl, and
15 ethoxycarbonyl, and R₄ is selected from the group consisting of methyl, o-nitrophenyl, and p-methoxyphenyl, and (d) R₂ is selected from the group consisting of hydrogen and methyl, R₃ and R₅ are same or different and selected from the group consisting of hydrogen,
20 methoxycarbonyl, and ethoxycarbonyl, R₄ is selected from the group consisting of methyl, o-nitrophenyl, and p-methoxyphenyl, and R₆ is selected from the group consisting of hydrogen, methyl, and phenyl.

The present invention further provides a compound of
25 the formula V



(V)

or a pharmaceutically acceptable salt thereof, wherein R₁
30 is selected from the group consisting of C₁-C₆ alkylcarbonyl, aryl C₁-C₆ alkylcarbonyl, aryl C₂-C₆

alkenylcarbonyl, C₁-C₆ alkyloxycarbonyl, amino C₁-C₆ alkylcarbonyl, and arylcarbonyl, wherein said aryl may be further substituted with halo, nitro, hydroxy, amino or cyano; and R₂ is hydrogen or halogen. Aryl includes
 5 phenyl, naphthyl, and aromatic moieties having 3 or 4 rings.


Particular embodiments include a compound of formula V wherein R₂ is hydrogen, and R₁ is ethylcarbonyl, benzoyl phenylethylcarbonyl, styrylcarbonyl, 4-
 10 nitrobenzylcarbonyl, 4-aminobenzylcarbonyl, or 3-iodo-4-aminobenzylcarbonyl.

"Aryl" in this application refers to phenyl, naphthyl, and aromatic groups with 3 or more rings, and preferably phenyl, unless otherwise described.

15 All of the aforesaid compounds of the present invention can be used as is or in the form of a pharmaceutical composition comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of the compound.

20 The present invention further provides a method of treating a mammal comprising selectively blocking one or more of the adenosine receptors of the mammal by administering to the mammal at least one compound of formulas I-V.

25 The present invention provides a method of treating a mammal comprising selectively blocking the adenosine receptors of a mammal by administering to said mammal at least one compound of the formula I



30


wherein R₁ and R₃ are selected from the group consisting of hydrogen, hydroxy, C₁-C₆ alkoxy, and C₁-C₆ alkylcarbonyloxy; R₂ is selected from the group consisting of hydroxy, C₁-C₆

alkoxy, and C₂-C₆ alkenoxy, said alkenoxy together with the carbon atom of the phenyl ring forming an oxygen heterocycle; R₄ is selected from the group consisting of phenyl, styryl, phenylbutadienyl, phenylacetylenyl, and -CH=N-phenyl, and substituted phenyl, styryl, phenylacetylenyl, and phenylbutadienyl, wherein the phenyl ring is substituted with 1 to 5 C₁-C₆ alkoxy groups; with the provisos that when R₃ is hydrogen, R₁ and R₂ are neither hydroxy nor alkoxy; when R₁, R₂, and R₃ are hydrogen, R₄ is neither phenyl nor alkyloxyphenyl; and when R₃ is hydrogen and R₄ is phenyl, neither R₁ nor R₂ is alkylcarbonyloxy.

Selected compounds of formula I are set forth in Tables 3-5. These compounds have affinity for adenosine receptors in general. Among these compounds, certain compounds have greater affinity for one type of adenosine receptors, e.g., A₁, than other types of adenosine receptors. Therefore these compounds can be used to selectively block that type of adenosine receptors for which they have greater affinity. Thus, for instance, compounds 3b, 4, 6, 7, 8, 10, 11b, 11c, 11d, 16, 20, 21 and 24 can be used to selectively block the A₁ adenosine receptors in a mammal.

The present invention further provides a method of treating a mammal comprising selectively blocking the adenosine receptors of a mammal by administering to said mammal at least one compound selected from the group consisting of genistein, (±)dihydrogenistein, sakuranetin, α-naphthoflavone, β-naphthoflavone, amaryllidaceae, oxogalanthine lactam, acetylhaemanthine methiodide, 2,3-methylenedioxy-fluorene-9-one, hematoxylin, and arborinine.

The present invention further provides a method of treating a mammal comprising selectively blocking one or more of the adenosine receptors of a mammal by administering to said mammal at least one compound of the formula II




wherein R_1 is selected from the group consisting of hydroxyl and C_1-C_6 alkoxy, and M is a divalent radical
5 selected from the group consisting of $-CH(OH)-CH(R_2)-$ and $-C(OH)=C(R_2)-$, wherein R_2 is selected from the group consisting of styryl and phenylacetylenyl.

Selected compounds of formula II are set forth in Tables 6-7. These compounds have affinity for adenosine
10 receptors in general. Among these compounds, certain compounds have greater affinity for one type of adenosine receptors, e.g., A_3 , than other types adenosine of receptors. Therefore these compounds can be used to selectively block that type of adenosine receptors for
15 which they have greater affinity. Thus, for instance, compound 38 can be used to selectively block the A_3 adenosine receptors in a mammal.

Selected compounds of formula III are set forth in Tables 10 and 18. These compounds have affinity for
20 adenosine receptors in general. Among these compounds, certain compounds have greater affinity for one type of receptors, e.g., A_1 , than other types adenosine of receptors. Therefore, these compounds can be used to selectively block that type of receptors for which they
25 have greater affinity. Thus, for instance, compounds 52, 58, 60, 61, 62, 63, 64, 65, 68, 69, 70, 71, 75, 76, and 77, which have greater affinity to A_1 adenosine receptors than A_{2a} receptors, can be used to selectively block A_1 adenosine receptors. Further, compounds 63, 64, 65, 74,
30 75, 76, 79, 87, 90, 93-95, 98-101, 105-107, 109, 115-126, 129-130b, 133, and 136, which have greater affinity for A_1 adenosine receptors than A_1 or A_{2a} adenosine receptors, can be used to selectively block A_3 adenosine receptors.

The present invention further provides a method of treating a mammal comprising selectively blocking one or more of the adenosine receptors of a mammal by administering to said mammal at least one compound of the
5 formula IV



or pharmaceutically acceptable salts thereof, wherein R_2 is selected from the group consisting of hydrogen and C_1-C_6 alkyl; R_3 is selected from the group consisting of hydrogen and C_1-C_6 alkyloxy; R_4 is selected from the group
10 consisting of C_1-C_6 alkyl, phenyl C_2-C_6 alkenyl, phenyl C_2-C_6 alkynyl, aryl, and aryl substituted with one or more substituents selected from the group consisting of nitro and C_1-C_6 alkyloxy; R_5 is selected from the group
15 consisting of hydrogen, C_1-C_6 alkyloxy carbonyl, and aryl C_1-C_6 alkyloxy carbonyl; and R_6 is selected from the group consisting of hydrogen, aryl, and C_1-C_6 alkyl.

The pharmaceutically acceptable carriers described herein, for example, vehicles, adjuvants, excipients, or
20 diluents, are well-known to those who are skilled in the art and are readily available to the public. It is preferred that the pharmaceutically acceptable carrier be one which is chemically inert to the active compounds and one which has no detrimental side effects or toxicity
25 under the conditions of use.

The choice of carrier will be determined in part by the particular active agent, as well as by the particular method used to administer the composition. Accordingly, there is a wide variety of suitable formulations of the
30 pharmaceutical composition of the present invention. The following formulations for oral, aerosol, parenteral, subcutaneous, intravenous, intraarterial, intramuscular, interperitoneal, intrathecal, rectal, and vaginal

administration are merely exemplary and are in no way limiting.

Formulations suitable for oral administration can consist of (a) liquid solutions, such as an effective
5 amount of the compound dissolved in diluents, such as water, saline, or orange juice; (b) capsules, sachets, tablets, lozenges, and troches, each containing a predetermined amount of the active ingredient, as solids or granules; (c) powders; (d) suspensions in an
10 appropriate liquid; and (e) suitable emulsions. Liquid formulations may include diluents, such as water and alcohols, for example, ethanol, benzyl alcohol, and the polyethylene alcohols, either with or without the addition of a pharmaceutically acceptable surfactant, suspending
15 agent, or emulsifying agent. Capsule forms can be of the ordinary hard- or soft-shelled gelatin type containing, for example, surfactants, lubricants, and inert fillers, such as lactose, sucrose, calcium phosphate, and corn starch. Tablet forms can include one or more of lactose,
20 sucrose, mannitol, corn starch, potato starch, alginic acid, microcrystalline cellulose, acacia, gelatin, guar gum, colloidal silicon dioxide, croscarmellose sodium, talc, magnesium stearate, calcium stearate, zinc stearate, stearic acid, and other excipients, colorants, diluents,
25 buffering agents, disintegrating agents, moistening agents, preservatives, flavoring agents, and pharmacologically compatible carriers. Lozenge forms can comprise the active ingredient in a flavor, usually sucrose and acacia or tragacanth, as well as pastilles
30 comprising the active ingredient in an inert base, such as gelatin and glycerin, or sucrose and acacia, emulsions, gels, and the like containing, in addition to the active ingredient, such carriers as are known in the art.

The compounds of the present invention, alone or in
35 combination with other suitable components, can be made into aerosol formulations to be administered via inhalation. These aerosol formulations can be placed into

pressurized acceptable propellants, such as dichlorodifluoromethane, propane, nitrogen, and the like. They also may be formulated as pharmaceuticals for non-pressured preparations, such as in a nebulizer or an atomizer.

Formulations suitable for parenteral administration include aqueous and non-aqueous, isotonic sterile injection solutions, which can contain anti-oxidants, buffers, bacteriostats, and solutes that render the formulation isotonic with the blood of the intended recipient, and aqueous and non-aqueous sterile suspensions that can include suspending agents, solubilizers, thickening agents, stabilizers, and preservatives. The compound can be administered in a physiologically acceptable diluent in a pharmaceutical carrier, such as a sterile liquid or mixture of liquids, including water, saline, aqueous dextrose and related sugar solutions, an alcohol, such as ethanol, isopropanol, or hexadecyl alcohol, glycols, such as propylene glycol or polyethylene glycol, glycerol ketals, such as 2,2-dimethyl-1,3-dioxolane-4-methanol, ethers, such as poly(ethyleneglycol) 400, an oil, a fatty acid, a fatty acid ester or glyceride, or an acetylated fatty acid glyceride with or without the addition of a pharmaceutically acceptable surfactant, such as a soap or a detergent, suspending agent, such as pectin, carbomers, methylcellulose, hydroxypropylmethylcellulose, or carboxymethylcellulose, or emulsifying agents and other pharmaceutical adjuvants.

Oils, which can be used in parenteral formulations include petroleum, animal, vegetable, or synthetic oils. Specific examples of oils include peanut, soybean, sesame, cottonseed, corn, olive, petrolatum, and mineral. Suitable fatty acids for use in parenteral formulations include oleic acid, stearic acid, and isostearic acid. Ethyl oleate and isopropyl myristate are examples of suitable fatty acid esters. Suitable soaps for use in parenteral formulations include fatty alkali metal,

ammonium, and triethanolamine salts, and suitable detergents include (a) cationic detergents such as, for example, dimethyl dialkyl ammonium halides, and alkyl pyridinium halides, (b) anionic detergents such as, for example, alkyl, aryl, and olefin sulfonates, alkyl, olefin, ether, and monoglyceride sulfates, and sulfosuccinates, (c) nonionic detergents such as, for example, fatty amine oxides, fatty acid alkanolamides, and polyoxyethylenepolypropylene copolymers, (d) amphoteric detergents such as, for example, alkyl- β -aminopropionates, and 2-alkyl-imidazoline quaternary ammonium salts, and (e) mixtures thereof.

The parenteral formulations will typically contain from about 0.5 to about 25% by weight of the active ingredient in solution. Suitable preservatives and buffers can be used in such formulations. In order to minimize or eliminate irritation at the site of injection, such compositions may contain one or more nonionic surfactants having a hydrophile-lipophile balance (HLB) of from about 12 to about 17. The quantity of surfactant in such formulations ranges from about 5 to about 15% by weight. Suitable surfactants include polyethylene sorbitan fatty acid esters, such as sorbitan monooleate and the high molecular weight adducts of ethylene oxide with a hydrophobic base, formed by the condensation of propylene oxide with propylene glycol. The parenteral formulations can be presented in unit-dose or multi-dose sealed containers, such as ampules and vials, and can be stored in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid carrier, for example, water, for injections, immediately prior to use.

Extemporaneous injection solutions and suspensions can be prepared from sterile powders, granules, and tablets of the kind previously described.

The compounds of the present invention may be made into injectable formulations. The requirements for effective pharmaceutical carriers for injectable

compositions are well known to those of ordinary skill in the art. See Pharmaceutics and Pharmacy Practice, J.B. Lippincott Co., Philadelphia, PA, Banker and Chalmers, eds., pages 238-250 (1982), and ASHP Handbook on
5 Injectable Drugs, Toissel, 4th ed., pages 622-630 (1986).

Additionally, the compounds of the present invention may be made into suppositories by mixing with a variety of bases, such as emulsifying bases or water-soluble bases. Formulations suitable for vaginal administration may be
10 presented as pessaries, tampons, creams, gels, pastes, foams, or spray formulas containing, in addition to the active ingredient, such carriers as are known in the art to be appropriate.

The compounds of the present invention can be used in
15 the treatment of any disease state or condition involving the release of inositol-1,4,5-triphosphate (IP3), diacylglycerol (DAG), and free radicals and subsequent arachidonic acid cascades. Thus, high blood pressure, locomotor hyperactivity, hypertension, acute hypoxia,
20 depression, and infertility can be treated in accordance with the present inventive method, wherein one of the above-described compounds is acutely administered, e.g., within about a few minutes to about an hour of the onset or realization of symptoms. The method also has utility
25 in the treatment of chronic disease states and conditions, in particular those conditions and disease states wherein chronic prophylactic or therapeutic administration of one of the above-described compounds will prevent the onset of symptoms or will reduce recovery time. Examples of
30 disease states and conditions that may be treated in accordance with the present inventive method include inflammatory disorders, such as vascular inflammation and arthritis, allergies, Crohn's disease, asthma, wound healing, stroke, cardiac failure, acute spinal cord
35 injury, acute head injury or trauma, seizure, neonatal hypoxia (cerebral palsy; prophylactic treatment involves chronic exposure through placental circulation), chronic

hypoxia due to arteriovenous malformations and occlusive cerebral artery disease, severe neurological disorders related to excitotoxicity, Parkinson's disease, Huntington's chorea, and other diseases of the central nervous system (CNS), cardiac disease, kidney disease, and contraception.

These compounds can be significant cerebral protectants. As such, the above compounds can be used to treat and/or protect against a variety of disorders, including, for example, seizures, transient ischemic shock, strokes, focal ischemia originating from thrombus or cerebral hemorrhage, global ischemia originating from cardiac arrest, trauma, neonatal palsy, hypovolemic shock, and hyperglycemia and associated neuropathies. The above method is applicable, for example, where a mammal has or is at risk of having a condition, disorder, or disease state associated with the cellular release of inositol-1,4,5-triphosphate or diacylglycerol. The method is also applicable when said mammal has or is at risk for hyperactivity and said compound in binding to said A₁ adenosine receptors functions as a locomotor depressant.

The present inventive method is also applicable when said mammal has or is at risk for hypertension and said compound in binding to said A₁ adenosine receptors functions as a hypotensive agent. The method is also applicable when said mammal has or is at risk for anxiety and said compound in binding to said A₁ adenosine receptors functions as an anxiolytic agent. The method is furthermore applicable when said mammal has or is at risk for cerebral ischemia and said compound in binding to said A₁ adenosine receptors functions as a cerebroprotectant. The method is also applicable when said mammal has or is at risk for seizures and said compound in binding to said A₁ adenosine receptors functions as an antiseizure agent.

The present inventive method can be administered chronically as well as acutely.

The present inventive method includes the administration to an animal, such as a mammal, particularly a human, in need of the desired adenosine receptor-dependent response of an effective amount, e.g.,
5 a therapeutically effective amount, of one or more of the aforementioned present inventive compounds or pharmaceutically acceptable salts or derivatives thereof, alone or in combination with one or more other pharmaceutically active compounds.

10 Any suitable pharmaceutically acceptable salts can be used. Example of suitable salts include carbonate, bicarbonate, sulfate, bisulfate, nitrate, halides, phosphates, oxalate, acetate, formate, citrates, and amino acid salts.

15 Some of the compounds of the present invention can be utilized as functionalized congeners for coupling to other molecules, such as amines and peptides. The use of such congeners provide for increased potency, prolonged duration of action, specificity of action, and prodrugs.
20 Water solubility is also enhanced, which allows for reduction, if not complete elimination, of undesirable binding to plasma proteins and partition into lipids. Accordingly, improved pharmacokinetics can be realized.

One skilled in the art will appreciate that suitable
25 methods of administering a compound of the present invention to an animal are available, and, although more than one route can be used to administer a particular compound, a particular route can provide a more immediate and more effective reaction than another route.
30 Accordingly, the above-described methods are merely exemplary and are in no way limiting.

The dose administered to an animal, particularly a human, in the context of the present invention should be sufficient to effect a prophylactic or other therapeutic
35 response in the animal over a reasonable time frame. One skilled in the art will recognize that dosage will depend upon a variety of factors including the strength of the

particular compound employed, the age, species, condition, and body weight of the animal, as well as the severity/stage of the disease or condition. The size of the dose will also be determined by the route, timing and frequency of administration as well as the existence, nature, and extent of any adverse side-effects that might accompany the administration of a particular compound and the desired physiological effect. It will be appreciated by one of skill in the art that various conditions or disease states, in particular chronic conditions or disease states, may require prolonged treatment involving multiple administrations.

Suitable doses and dosage regimens can be determined by conventional range-finding techniques known to those of ordinary skill in the art. Generally, treatment is initiated with smaller dosages, which are less than the optimum dose of the compound. Thereafter, the dosage is increased by small increments until the optimum effect under the circumstances is reached. For convenience, the total daily dosage may be divided and administered in portions during the day if desired. In proper doses and with suitable administration of certain compounds, the present invention provides for a wide range of selective adenosine receptor-dependent responses. Exemplary dosages range from about 0.01 to about 100 mg/kg body weight of the animal being treated/day. Preferred dosages range from about 0.1 to about 10 mg/kg body weight/day.

The abbreviations used in this application have the following meaning:

30	[¹²⁵ I]AB-MECA	[¹²⁵ I]N6-(4-amino-3-iodobenzyl)adenosine-5'-N-methyluronamide
	CHO	Chinese hamster ovary
	DMF	N,N-dimethylformamide
	DMSO	dimethylsulfoxide
35	K _i	equilibrium inhibition constant
	R-PIA	R-N6-phenylisopropyladenosine
	Bay K 8422:	

1,4-dihydro-2,6-dimethyl-5-nitro-4-[2-(trifluoromethyl)-phenyl]-3-pyridine carboxylic acid methyl ester

The following examples further illustrate the present invention but, of course, should not be construed as in
5 any way limiting its scope.

EXAMPLE 1

This Example illustrates the various analytical methods employed in the characterization of the compounds
10 of the present invention.

Proton nuclear magnetic resonance spectroscopy was performed on a Varian GEMINI-300 spectrometer and spectra were taken in d6-DMSO. Electron-impact mass spectrometry was performed with a VG7070F mass spectrometer at 6 kV.
15 Elemental analysis was performed by Atlantic Microlabs, Inc. (Norcross, GA).

EXAMPLE 2

This Example illustrates the procedure for
20 determining the affinity of the present inventive compounds for adenosine receptors.

K_i values at A_1 and A_{2a} receptors were determined in radioligand binding assays in brain membranes vs. [3H]PIA or [3H]CGS 21680, respectively. Schwabe et al.,
25 Naunyn-Schmiedeberg's Arch. Pharmacol., **313**, 179-187 (1980); Jarvis et al., J. Pharmacol. Exp. Therap., **251**, 888-893 (1989). Affinity at human brain A_1 receptors expressed in HEK-293 cells, Salvatore et al., Proc. Natl. Acad. Sci., **90**, pp. 10365-10369 (1993), was determined
30 using [125 I]AB-MECA. Olah et al., Mol. Pharmacol., **45**, 978-982 (1994). Compounds were tested at human A_1 receptors, and selected compounds, which would be preferred for comparison with the A_1 and A_{2a} data, were also tested at rat A_1 receptors stably expressed in Chinese
35 hamster ovary (CHO) cells for inhibition of adenylyl cyclase. Kim, H.O., J. Med. Chem., **37**, 3614-3621 (1994).

The human A_1 receptor was chosen for several reasons

including that the affinity of most known adenosine adenosine ligands is minimal at rat A₁ receptors. In addition, the human A₁ receptor, being more sensitive, allowed for a better comparison between compounds.

- 5 Salvatore et al., Proc. Natl. Acad. Sci., 90, 10365-10369 (1993).

EXAMPLE 3

10 This Example illustrates the general synthetic methodology of certain flavone derivatives of the present invention.

The synthesis of the flavone derivatives can be accomplished following the teachings in literature and the procedures set forth herein. Wollenweber et al., J.B. Harborne, (Chapman and Hall, eds.), London, 259-335 (1993); Cushman et al., J. Med. Chem., 37, 3373-3382 (1994); Cunningham et al., Anti-Cancer Drug Design, 7, 365-384 (1992); Lee et al., Tetrahedron, 51, 4909-4922 (1995). As shown in Figure 2, the flavone ring system could be formed via a condensation of a 2-hydroxyacetophenone derivative, Lee et al., Tetrahedron, 51, 4909-4922 (1995), with the appropriate aldehyde, Williams, Spectroscopic Methods in Organic Chemistry, 4th ed., McGraw Hill, London, 1989, which gives rise to the 2-substituent. A cinnamaldehyde derivative was used as the second component thus leading to cinnamyl substitution at the 2-position. The condensation resulted first in a trans-olefin, Duarte et al., Eur. J. Pharmacol., 286, 115-122 (1995), which could be cyclized under basic oxidizing conditions to give either the 2,3-saturated dihydroflavonol analogues, Kim et al., J. Med. Chem., 37, 3373-3382 (1994), and Siddiqi et al., Nucleosides Nucleotides (1995, in press), or at higher temperature the corresponding dehydro derivative, Palmer et al., J. Biol. Chem., 270, 16895-16902 (1995), a flavonol. Evidence that the 3-hydroxyflavanones obtained were of the trans configuration is from NMR. The coupling constant for the

2- and 3-position protons for compounds 15 - 16 is ~12 Hz, which is characteristic of trans vicinal coupling.

Williams et al., Spectroscopic Methods in Organic Chemistry, 4th ed., McGraw Hill, London (1989).

5 An alternate approach to providing olefinic substitution at the 2-position (Figure 2) was to condense a 2-methylchromone (such as the natural product visnagin, 17)27 with an aldehyde, Hafez, Czech. Chem. Commun., 59, (1994), under basic conditions. This method was used to
10 prepare the 2-cinnamyl derivative, Ji et al., J. Med. Chem. (1996, in press). Evidence for a trans-styryl group of Ji et al., J. Med. Chem. (1996, in press), was found in the NMR spectra with the olefinic coupling constant of 15 Hz. In the presence of ethoxide ions the 5-methoxy group
15 readily exchanged for ethoxy, resulting in compound 20. The corresponding 5-propoxy derivative, 21, Markham et al., J.B. Harborne, (Chapman and Hall eds., London, 441-497 (1993), was obtained using an alternate approach in which visnagin was demethylated to give Schwabe et al.,
20 Naunyn-Schmiedeberg's Arch. Pharmacol., 313, 179-187 (1980), followed by condensation with benzaldehyde. The 2-methyl group of visnagin could also be oxidized to an aldehyde group, as in 18. Hafez M. El-Shaaer, Collect. Czech. Chem. Commun., 59 (1994).

25

EXAMPLE 4

This Example illustrates the alkylation of galangin, 1 and morin, 5a.

30 The alkylation of galangin was carried out using the appropriate alkyl iodide and potassium carbonate in refluxing acetone (Figure 1), providing the completely alkylated product, 3a or 4. Partial alkylation leaving the 5-position hydroxyl group unreacted, to provide 3b, was accomplished under less forceful conditions, e.g.
35 alkyl bromide at room temperature. The O-alkylation of morin, 5a, (2',3,4',5,7-pentahydroxyflavone) was complicated by an even lesser reactivity of the 5-hydroxy

group than in galangin, leading to tetra- rather than penta- substitution (Figure 1). Thus, when ethyl sulfate was used the fully alkylated morin derivative, 8, was obtained, while reaction with ethyl iodide provided only the tetraethyl derivative, 7. The position of the non-alkylated hydroxyl group of 3b and 7 was determined using NMR. The resonance for the 5-OH proton appears at roughly 12 ppm. Kim et al., *J. Med. Chem.*, **37**, 3614-3621 (1994). The downfield shift is apparently caused by the adjacent carbonyl to which it may H-bonding, forming a six-membered ring. This proton signal is present in the spectra of 3b, 5a and 7 but not 8.

The physical characterization data of the flavonoid derivatives described in this Example as well as in the following Examples are set forth in Tables 1 and 2.

3,5,7-Trimethoxyflavone (2)

Galangin (27 mg, 0.1 mmol) was dissolved in dried acetone (20 mL), solid potassium carbonate (0.5 g) and dimethyl sulfate (1 mL) were added, and the mixture was refluxed for 4 hrs, then cooled to room temperature. The solution was filtered and evaporated, water (20 mL) and concentrated ammonium hydroxide (2 mL) were added and the solution was extracted with ethyl acetate (15 mL x 2). The solvent was evaporated and the residue was recrystallized from methanol/water to give 21 mg product (67%). ¹HNMR (DMSO-d₆): δ 3.73 (s, 3H, 3-OCH₃), 3.86 (s, 3H, 7-OCH₃), 3.86 (s, 3H, 5-OCH₃), 6.48 (s, 1H, 8-H), 6.79 (s, 1H, 6-H), 7.99 (m, 3H), 8.02 (m, 2H). MS (CI/NH₃): m/z 313 (MH⁺, base).

3,5,7-Triethoxyflavone (3a)

Galangin (30 mg, 0.11 mmole) was dissolved in dried acetone (45 mL), solid potassium carbonate (0.5 g) and iodoethane were added, and the mixture was refluxed for overnight. The solution was filtered and evaporated, water (20 mL) and concentrated ammonia (1 mL) were added,

and the solution was extracted with ethyl acetate. The ethyl acetate was evaporated, and the crude mass was purified on a preparative silica TLC plate to give 12 mg 3,5,7 triethoxyflavone. mp 111-114°C. mass (FAB) m/z 355 (M++1, base) 1H NMR (CDCl₃) 1.33 (t, 3H, CH₃), 1.55 (t, 3H, CH₃), 1.62 (t, 3H, CH₃), 4.1-4.25 (m, 6H, CH₂), 6.4 (s, 1H), 6.55 (s, 1H), 7.5-7.6 (m, 3H), 8.1-8.2.

3, 7-Diethoxy-5-hydroxyflavone (3b)

Galangin (27 mg, 0.1 mmol) was dissolved in dried acetone (20 mL), solid potassium carbonate (0.5 g) and bromoethane (1 mL) were added and the mixture was stirred overnight, at room temperature. The solution was filtered and evaporated, water (20 mL) and concentrated ammonium hydroxide (2 mL) were added and the solution was extracted with ethyl acetate (20 mL x 2). The solution was dried and the solvent was evaporated. The residue was purified by preparative TLC plate (silica, ethyl acetate/petroleum ether 2 : 8) to give 21 mg product (64%). 1HNMR (CDCl₃): d 1.34 (t, 3H, J = 7.3 Hz, 3-CH₃), 1.48 (t, 3H, J = 7.1 Hz, 7-CH₃), 4.14 (m, 4H, 2 x OCH₂), 6.38 (s, 1H, 6-H), 6.47 (s, 1H, 8-H), 12.64 (s, 1H, 5-OH). MS (CI/NH₃): m/z 327 (MH⁺, base).

3,5,7-Tripropyloxyflavone (4)

Galangin (30 mg, 0.11 mmole) was dissolved in dried acetone (45 mL), solid potassium carbonate (0.5 g) and 1-iodopropane were added and the mixture was refluxed overnight. The solution was filtered and evaporated, water (20 mL) and concentrated ammonia (1 mL) were added, and the solution was extracted with ethyl acetate. The ethyl acetate was evaporated, and the crude mass was purified on a preparative TLC plate (silica) to give 25 mg (63%) of 3,5,7-tripropyloxyflavone, mp 90-93°C. Mass spectra (CI-NH₃) m/z 397 (M++1), base) 1H NMR (CDCl₃)

0.85-1.2 (m, 9H), 1.6-2.1 (m, 6H), 3.9-4.1 (m, 6H), 6.32 (s, 1H), 6.5 (s, 1H), 7.4-7.5 (m, 3H), 8.04-8.12 (m, 2H).

2',3,4',5,7-Pentamethoxyflavone (6)

5 Morin (120 mg, 0.4 mmol) was dissolved in dried acetone (80 mL). Solid potassium carbonate (2 g) and iodomethane (2 mL) were added, and the mixture was refluxed overnight, then cooled to room temperature. The solution was filtered and evaporated, and the residue was
10 separated by preparative TLC plates (silica, ethyl acetate) to give 80 mg product (55%). ¹H NMR (CDCl₃): δ 3.86, 3.88, 3.91, 3.93, 4.02 (5s, 15H, 5 x OCH₃), 6.39, 6.49, 6.62 (s, 1H, 8-H), 6.79 (3m, 3H, phenyl), 6.65 (m, 1H, 8-H), 7.42 (m, 1H, 6-H)). MS (EI): m/z 372 (M⁺), 371
15 (M-H)⁺, 341 (M-OCH₃)⁺.

2',4',3,7-Tetraethoxy-5-hydroxymorin (7)

Compound 7 was synthesized according to the above procedure for 6, except for using iodoethane instead of
20 iodomethane. The product, 7, was separated on preparative TLC with petroleum ether/ethyl acetate and displayed a mass (FAB) m/z 415 (M⁺⁺¹, base). ¹H NMR (CDCl₃) 1.15 (t, 3H, CH₃), 1.35 (t, 3H, CH₃), 1.45 (dt, 6H, CH₃), 3.9-4.2 (m, 8H, CH₂), 6.33 (d, 1H), 6.6 (m, 3H), 7.4 (d, 1H).

2',3,4',5,7-Pentaethoxyflavone (8)

Morin (302 mg, 1 mmol) was dissolved in dried acetone (80 mL), solid potassium carbonate (5 g) and diethyl sulfate (5 mL) were added, and the mixture was refluxed overnight, then cooled to room temperature. The solution was filtered and evaporated, water (50 mL) and concentrated ammonium hydroxide (10 mL) were added and the solution was extracted with ethyl acetate (40 mL x 2). The residue was crystallized from ethyl acetate and petroleum ether (1 : 9) to give 335 mg product (76%). ¹H NMR (CDCl₃): d 1.12 (t, 3H, J = 7.1 Hz, CH₃), 1.32 (t, 3H, J = 7.1 Hz, CH₃), 1.45 (2t, overlap, 6H, 2 x CH₃), 1.56 (t, 3H, J = 7.1 Hz, CH₃), 4.10 (m, 10 H, 5 x OCH₂), 6.32 (d, 1H, J = 1.2 Hz, 2-phenyl), 6.39 (s, 1H, 2-phenyl), 6.53 (s, 1H, 2-phenyl), 6.58 (s, 1H, 6-H), 7.37 (d, 1H, 8-H). MS (CI/NH₃): m/z 443 (MH⁺, base).

EXAMPLE 5

This Example illustrates the synthesis of 2-phenylacetylenyl-3-hydroxy-6-methoxyflavone (10).

1-(2-Hydroxy-5-methoxy)-5-phenylpenta-2-en-4-yne-1-one (Compound 15a, 60 mg, 0.2 mmole) was dissolved in ethanol (1.5 mL). Sodium hydroxide 1N (0.4 mL) and hydrogen peroxide (0.5g) were added, and the mixture was heated at 90°C for 35 min. The cooled mixture was diluted with water (5 mL) and acidified with hydrochloric acid, 1N. A precipitate was collected and recrystallized from methanol. Mass (CI NH₃) m/z 293 (M⁺⁺¹, base), ¹H NMR (CDCl₃) 3.95 (s, 3H, CH₃), 6.92 (s, 1H), 7.25-8.0 (m, 7H).

EXAMPLE 6

This Example illustrates the synthesis of trans-2-Styryl-3-hydroxy-6-methoxyflavanone (15).

2'-Hydroxy-5'-methoxyacetophenone (4.1 g, 22 mmole) and cinnamaldehyde (3.35 g, 25 mmole) were dissolved in minimum methanol (2.5 mL). Concentrated sodium hydroxide

(12.5 mL, 50%) was added, and the mixture was kept on ice for 8 hr. The resultant solid was suspended in water and acidified using HCl (4N). The oil that separated was dissolved in ethanol and crystallized from ethanol/water to give

1- (2-hydroxy-5-methylphenyl)-5-phenylpenta-2,4-dien-1-one (1.2 g) as red-brown powder. Mass (CI-NH₃) m/z 281 (M++1, base). ¹H NMR (CDCl₃) .3.9(s, 3H, CH₃), 6.95-7.8(m, 12H). The above compound (170 mg, 0.6 mmole) was dissolved in a mixture of ethanol (3.5mL) and acetone (4mL). Sodium hydroxide 1N (1 mL) and hydrogen peroxide (1 mL, 35%) were added, and the solution was stirred 6 hr. at room temperature. The mixture was precipitated by adding water and HCl and purified on preparative TLC (petroleum ether-ethyl acetate) to give compound 15, mp 160-163°C. mass (CI NH₃) m/z 297(M++1,base). ¹H NMR (CDCl₃) 3.9(s, 3H, CH₃), 4.45(d, J=12Hz, 1H), 4.8(m, 1H), 6.55(dd, 1H), 6.9-7.6(m, 9H).

20

EXAMPLE 7

This Example illustrates the synthesis of trans-2-phenylacetylenyl-3-hydroxy-6-methoxyflavanone (16).

2'-Hydroxy-5'-methoxyacetophenone and phenylphenylacetylenyl aldehyde were condensed according to the procedure for the preparation of compound 15 to give 1- (2-hydroxy-5-methylphenyl)-5-phenylpenta-2-en-4-yne-1-one. Mass (CI-NH₃) m/z 279 (M++1, base). ¹H NMR (CDCl₃) .3.9 (s, 3H, CH₃), 7.0 (d, 1H), 7.1-7.6 (m, 9H). The above compound was allowed to react with hydrogen peroxide to give compound 16 as a white powder, mp 135-138°C. Mass (EI) m/z 295 (M+1) 150 (base). ¹H NMR (CDCl₃) d 3.85 (s, 3H, OCH₃), 4.6 (d, J=12Hz, 1H), 5.09 (d, J=12Hz, 1H), 7.0-7.6 (m, 8H). 4-Methoxy-7-formyl-5H-furo[3,2-g][1]-benzopyrano-5-one (18) was synthesized from visnagin (4-methoxy-7-methyl-5H-furo[3,2-g][1]-

benzopyrano-5-one) according to Duarte et al., Eur. J. Pharmacol., 286, 115-122 (1995).

EXAMPLE 8

5 This Example illustrates the synthesis of 4-methyloxy-7-trans-styryl-visnagin (19) and 4-ethyloxy-7-trans-styryl-visnagin (20).

10 Visnagin (17, 160 mg, 0.7 mmole) and benzaldehyde (120 mg, 1.1 mmole) were dissolved in ethanol (4 mL). Sodium ethoxide (20% in ethanol, 0.5 ml) was added, and the mixture was stirred for 10 min at 80°C. Disappearance of the starting material was accompanied with formation of two products Rf 0.7 and 0.8 in ethyl acetate. Both products were separated on preparative TLC plates (ethyl acetate) to give compound 19, mp 175-178°C. Mass (CI NH₃) m/z 319 (M++1, base) 1H NMR (CDCl₃) δ 4.2 (s, 3H, OCH₃), 6.23 (s, 1H), 6.75 (d, J=15Hz, 1H), 7.05 (s, 1H), 7.3-7.8 (m, 8H).

15 Compound 20, mp 148-151°C. Mass (CI NH₃) m/z 333 (M++1, base). 1H NMR (CDCl₃) 1.55 (t, 3H, CH₃), 4.4 (q, 2H, OCH₂), 6.23 (s, 1H), 6.78 (d, J=15Hz, 1H), 7.0 (s, 1H), 7.35-7.7 (m, 8H).

EXAMPLE 9

25 This Example illustrates the synthesis of 4-propyloxy-7-trans-styryl-visnagin (21).

A mixture of visnagin (2 g, 8.7 mmol), potassium iodide (10 g) and acetic acid (50 mL) was refluxed for 7 hrs. After cooling, the precipitate was removed by 30 filtration, and the filtrate was evaporated under reduced pressure, and coevaporated with toluene (10 mL x2). The residue was crystallized from ethanol to give 1.62 g (81%) of demethylated product of visnagin, 17a. 1HNMR (CDCl₃): δ 2.40 (s, 3H, 7-CH₃), 6.12 (s, 1H, 6-H), 7.00 and 7.62 (2d, 2H, J = 2.9 Hz 3-H and 2H), 7.05 (s, 1H, 9-H), 13.6 (s, 35 1H, 4-OH). The mixture of demethylated visnagin (430 mg,

2 mmol), iodopropane (3 mL) and potassium carbonate (5 g) in dry acetone (80 mL) was refluxed overnight. The solid was removed by filtration, and the solvent was evaporated, water (50 mL) and concentrated ammonium hydroxide (15 mL) were added, and the solution was extracted with ethyl acetate (40 mL x 2). The organic layer was dried over sodium sulfate, and the solvent was evaporated. The residue was purified by preparative TLC plates (silica, ethyl acetate/petroleum ether, 4:6) to give 465 mg (90%) of 4-propyloxy-visnagin, **17b**. ¹HNMR (CDCl₃): δ 1.10 (t, 3H, J = 7.5 Hz), 1.94 (m, 2H), 2.33 (s, 3H, 7-CH₃), 4.23 (t, 2H, J = 6.7 Hz, 4-OCH₂), 6.03 (s, 1H, 6-H), 6.97 and 7.60 (2d, 2H, J = 2.7 Hz, 3-H and 2H), 7.27 (s, 1H, 9-H).

4-propyloxy-visnagin (86 mg, 0.3 mmol) and benzaldehyde (50 mg, 0.5 mmol) were dissolved in ethanol (2 mL). While stirring, sodium ethoxide (20% in ethanol, 0.3 mL) was added, and the mixture was stirred for 15 min at 80 °C. The solvent was evaporated, and the residue was partitioned between ice water (20 mL) and ethyl acetate (20 mL), the aqueous layer was extracted with ethyl acetate (20 mL), and the combined organic layer was dried over sodium sulfate. The solvent was evaporated, and the residue was separated by preparative TLC plate (silica, ethyl acetate/petroleum ether, 4:6) to give 26 mg of the desired product, **21** (25% yield). ¹HNMR (CDCl₃): δ 1.07 (t, 3H, J = 7.5 Hz), 1.97 (m, 2H), 4.25 (t, 2H, J = 6.7 Hz, 4-OCH₂), 6.21 (s, 1H, 6-H), 6.78 (d, 2H, J = 14 Hz, 3-H), 7.00 (s, 1H, 9-H), 7.33 (m, 8H). MS (CI/NH₃): m/z 347 (MH⁺, base).

30

EXAMPLE 10

This Example illustrates the synthesis of 4-methoxy-7-phenylbutadienyl visnagin (**22**) and 4-ethoxy-7-phenylbutadienyl visnagin (**23**).

Compound **22** was prepared by dissolving visnagin and cinnamaldehyde in ethanol in the presence of sodium ethoxide according to the above procedure for preparing

compounds 19 and 20 to yield compound 22, mp 162-165°C.
Mass (EI) m/z 344 (M+, base). ¹H NMR (CDCl₃) δ 4.2 (s, 3H, OCH₃), 6.15 (s, 1H), 6.33 (d, J=15 Hz, 1H), 6.95 (s, 1H), 7.0-7.6 (m, 10H).

- 5 Compound 23 was also isolated from the above reaction. Mass (EI) m/z 358 (M+), 343 (M-15, base). ¹H NMR (CDCl₃) 1.55 (t, 3H, CH₃), 4.4 (q, 2H, OCH₂), 6.15 (s, 1H), 6.35 (d, J=15Hz, 1H), 7.4 (d, J=15Hz, 1H), 6.9-7.6 (m, 10H).

10

EXAMPLE 11

This Example illustrates the synthesis of compound 24 (the Schiff base of compound 18).

- Compound 18 (400 mg) and aniline (0.5 mL) were
15 dissolved in toluene and stirred overnight at 25°C. The course of the reaction was followed using analytical TLC (silica, CHCl₃/MeOH, 20:1). The solvent was evaporated, and excess aniline was removed under high vacuum. The product (R_f ~ 0.7) was purified using preparative TLC
20 (silica, ethyl acetate). Mass (CI NH₃) m/z 320 (M++1, base).

EXAMPLE 12

- This Example illustrates the synthesis of
25 3,5,7-triacetoxyflavone (34).

- Galangin (9.0 mg, 33 μmol) was dissolved in 1 mL DMF and treated with acetic anhydride (0.2 mL) and 4,4-dimethylaminopyridine (3 mg). After stirring for 10 min, 2 mL of 1 N NaH₂PO₄ was added. A white precipitate
30 was removed by filtration, and recrystallized from methanol/water to yield 11.4 mg of a solid (87%), which was homogeneous by TLC (R_f 0.56, silica, chloroform: methanol: acetic acid, 95:4:1). Mass spec: 414 (m+1+NH₃), 397 (m+1), 355 (m+1+NH₃-OAc). Mp 128-129 °C.
35 NMR δ 7.6 (3H, Ar), 7.9 (m, 2H, Ar), 7.65 and 7.17 (each,

d, 1H, J = 2 Hz, Ar, o- to AcO), 2.34 (s, 6H, Ac), 2.30 (s, 3H, Ac). CHN analysis.

EXAMPLE 13

5 This Example illustrates another method of synthesis of 2',3,4',5,7-pentamethoxyflavone (38, also designated as compound 6 in Example 4).

Morin hydrate (25 mg, 82 μ mol, Aldrich) was dissolved in dry acetone (20 mL), in which was suspended potassium
10 carbonate (1.0 g). Dimethyl sulfate (1.0 mL, 11 mmol) was added, and the mixture was refluxed for four h under nitrogen. After cooling in an ice bath, 2 mL concentrated ammonium hydroxide were added in aliquots followed by 20 mL water. The solution was extracted with ethyl acetate.

15 The organic layer was dried and evaporated, and the white solid residue was recrystallized from methanol/water to yield 11 mg (36%) of 20, which was homogeneous by thin layer chromatography (Rf 0.32, silica, chloroform:methanol:acetic acid, 95:4:1). Mass spec: 373
20 (m+1), 359 (m+1-CH₂), 343. Mp. 153-156 °C. NMR d 7.35 (d, 1H, J = 8 Hz, Ar, 6-Ph), 6.66 (dd, 1H, J = 2, 8 Hz, Ar, 5-Ph), 6.70 and 6.62 and 6.48 (each, d, 1H, J = 2 Hz, Ar), 3.84 (s, 9H, Me), 3.80 (s, 3H, Me), 3.61 (s, 3H, Me). CHN analysis.

25

EXAMPLE 14

This Example illustrates the synthesis of compound 45.

Compound 45 was synthesized by a modification of a
30 literature procedure. Fales et al. J. Amer. Chem. Soc., 77, 5885-5890 (1955). 6-Bromo-3,4-dimethoxybenzoic acid (6-bromoveratric acid, 2.0 g, 7.7 mmol, Spectrum Chem. Corp., New Brunswick, NJ) was dissolved in 50% EtOH/H₂O (100 mL), and treated with resorcinol (0.85 g, 7.7 mmol),
35 50 mg copper powder, 50 mg cupric acetate, and sodium hydroxide (0.31 g, 7.7 mmol). The mixture was heated to reflux overnight. The reaction mixture was extracted once

with ether, and insoluble solids were removed by filtration. The aqueous layer was acidified with 1 N HCl to form the lactone. The precipitate was removed by filtration and dried. In order to remove unreacted 6-bromoveratric acid from this solid containing the product, it was suspended and partially dissolved in cold saturated aqueous NaHCO₃. The remaining insoluble solids were collected by filtration and the washings discarded. The dried residue was recrystallized from EtOH to give 0.74 g of the pure product (35% yield). NMR: CDCl₃ δ ppm 8.68 and 7.65 (each s, 1H, Ar), 4.06 and 3.99 (each s, 3H, Me), 2.95 and 2.67 (each t, 2H, CH₂), 2.18 (m, 2H, CH₂). Mp. 236-237 °C. CHN analysis.

15

EXAMPLE 15

This Example illustrates the synthesis of 2,3-methylenedioxy-fluorene-9-one (49).

Gaseous HF was condensed by collecting in a Teflon vial placed in a dry ice/ acetone bath.

6-Phenylpiperonylic acid (100 mg, 0.41 mmol) was added to HF (2 mL) and the solution let stand overnight. The HF was evaporated leaving the pure fluorenone derivative in quantitative yield (92 mg), mp 156-157 °C. CHN analysis was carried out.

Table 1. Physical characterization of flavone, flavanol, and flavanone derivatives.

Compound	Formula	m.p. (°C)	M.W. (anhyd)	M.W. (MS) ^a	Elemental Analysis	
					Calculated	Found
2	C ₁₃ H ₁₂ O ₆ -0.75H ₂ O	201	336.4	313 (CI)	C, 66.35; H, 5.41	C, 66.39; H, 5.27
3a	C ₂₁ H ₁₆ O ₆	111-114	364.4	355 (FAB)	C, 71.17; H, 6.26	C, 70.94; H, 6.37
3b	C ₁₉ H ₁₂ O ₆	100-101	336.4	337 (CI)	C, 69.92; H, 5.56	C, 69.67; H, 5.66
4	C ₂₃ H ₂₀ O ₆	90-93	392.4	397 (CI)	C, 72.70; H, 7.12	C, 72.61; H, 7.14
6	C ₂₀ H ₂₀ O ₇	153-156	372.4	373 (CI)	C, 64.51; H, 5.41	C, 64.42; H, 5.44
7	C ₂₃ H ₂₆ O ₇	122-123		415 (FAB)		
8	C ₂₅ H ₃₀ O ₇	115	442.5	443 (CI)	C, 67.85; H, 6.83	C, 67.75; H, 6.84
10	C ₁₈ H ₁₂ O ₇	oil		293 (CI)		
15	C ₁₈ H ₁₆ O ₄ -0.25H ₂ O	160-163	296.3	297 (CI)	C, 71.86 H, 5.53	C, 71.93; H, 5.61
16	C ₁₈ H ₁₄ O ₄	135-138	294.3	294 (EI)	C, 73.46; H, 4.80	C, 74.30; H, 5.27
19	C ₂₀ H ₁₄ O ₄ -0.25H ₂ O	175-178	318.3	319 (CI)	C, 74.41; H, 4.53	C, 74.55; H, 4.47
20	C ₂₁ H ₁₆ O ₄	148-151	332.4	333 (CI)	C, 75.89; H, 4.85	C, 75.83; H, 4.86

Compound	Formula	m.p. (°C)	M.W. (anhyd)	M.W. (MS) ^a	Elemental Analysis	
					Calculated	Found
21	C ₂₂ H ₁₈ O ₄	132	346.4	347 (CI)	C, 76.30; H, 5.20	C, 76.03; H, 5.28
22	C ₂₂ H ₁₆ O ₄ ·0.5H ₂ O	162-165	344.4	344 (EI)	C, 74.77; H, 4.85	C, 74.98; H, 4.77
23	C ₂₃ H ₁₈ O ₄	162-165	358.4	358 (EI)	C, 77.08; H, 5.06	C, 76.91; H, 5.16
24	C ₁₉ H ₁₄ NO ₄	150-154	320.3	320 (CI)	C, 71.24; H, 4.41	N, 4.37; C, 71.50; H, 4.31; N, 4.37

a. High resolution mass in FAB+ mode m/z determined to be within acceptable limits.


Table 2. Elemental Analysis of Flavonoid Derivatives 36, 37, 41, 47, and 51.

Comp. No.	Formula	MW (anhydrous)	Calculated	Found
36	$C_{19}H_{12}O_6 \cdot 0.75H_2O$	336.4	C, 66.35; H, 5.41	C, 66.39; H, 5.27
37	$C_{21}H_{16}O_8 \cdot 0.25H_2O$	396.4	C, 62.92; H, 4.15	C, 62.76; H, 4.14
41	$C_{20}H_{20}O_7 \cdot 0.25H_2O$	372.4	C, 63.74; H, 5.48	C, 63.87; H, 5.80
47	$C_{13}H_{14}O_5$	274.3	C, 65.68; H, 6.52	C, 65.44; H, 5.20
51	$C_{14}H_8O_3 \cdot 0.1H_2O$	224.2	C, 74.39; H, 3.66	C, 74.33; H, 3.81

EXAMPLE 16

This Example illustrates the efficacy of some the flavonoid derivatives of the present invention in blocking adenosine receptors. The affinities of the flavonoid derivatives 1-11d were determined using radioligand binding assays, and the results thereof are set forth in Table 3.

Table 3. Affinities of flavonoid derivatives in radioligand binding assays at A_1 , A_{2a} , and A_3 receptors.^{a-e}



K_i (μ M) or % inhibition^c

Compound	R_1, R_2	R_3	R_4	rA_1^a	rA_{2a}^b	hA_3^c	rA_1/hA_3
1 (galangin)	5,7-(OH) ₂	OH	Ph	0.863 ± 0.092	0.966 \pm 0.164	3.15 \pm 0.85	0.27
2	5,7-(OMe) ₂	OMe	Ph	0.509 ± 0.049	6.45 \pm 1.48	1.21 \pm 0.30	0.42
3a	5,7-(OEt) ₂	OEt	Ph	0.603 ± 0.022	3.31 \pm 1.24	0.364 \pm 0.067	1.7
3b	5-OH, 7-OEt	OEt	Ph	1.89 \pm 0.56	63.8 \pm 18.8	0.748 \pm 0.402	2.5
4	5,7-(OPr) ₂	OPr	Ph	1.10 \pm 0.20	3.22 \pm 1.31	0.317 \pm 0.089	3.5
5a morin	5,7-(OH) ₂	OH	2',4'-(OH) ₂ - Ph	13.8 \pm 3.1	17.3 \pm 4.1	34 \pm 4% (10^{-4})	~ 0.1
5b	5-OH-6- CH=CH- CH(CH ₃) ₂ -	CH ₂ CH=CH- CH(CH ₃) ₂ -	2',4'-(OH) ₂ - Ph	9.09 \pm 0.91	d (10^{-4})	4.59 \pm 1.69	2.0


Compound	R ₁ , R ₂	R ₃	R ₄	rA ^a ₁	rA ^b _{2a}	hA ^c ₃	rA ₁ /hA ₃
7OMe							
6	5,7-(OMe) ₂	OMe	2',4'-(OMe) ₂ -Ph	27.6±7.5	46.7±2.7	2.65±0.72	10
7	5OH-7-OEt	OEt	2',4'-(OEt) ₂ -Ph	d (10 ⁻⁴)	d (10 ⁻⁴)	4.83±1.40	>40
8	5,7-(OEt) ₂	OEt	2',4'-(OEt) ₂ -Ph	32.3±9.1	26.8	7.27±1.88	4.4
9	H	OH	2',4',6'-(OMe) ₃ -Ph	7.32±1.36	d (10 ⁻⁴)	50.1±7.8	0.15
10	6-OCH ₃	OH	C≡C-Ph	d (10 ⁻⁴)	d (10 ⁻⁴)	24.0±9.7	>8
11a flavone	H	H	Ph	3.28±0.92	3.45±1.16	16.9±3.8	0.19
11b ^f	d	Cl	Ph	15% (10 ⁻⁴)	54.5±26.3	0.291±0.042	>300
11c ^f	d	Cl	2'-i-PrO-4'-Me-Ph	36±12% (10 ⁻⁴)	19±9% (10 ⁻⁴)	0.561±0.129	~200
11d ^f	d	Cl	2',4',6'-Me ₃ -Ph	23% (10 ⁻⁴)	d (10 ⁻⁴)	5.24±0.52	~20

- a Displacement of specific [³H]PIA binding in rat brain membranes, expressed as $K_i \pm$ S.E.M. in μ M (n = 3-5), or as a percentage of specific binding displaced at 100 μ M.
- b Displacement of specific [³H]CGS 21680 binding in rat striatal membranes, expressed as $K_i \pm$ S.E.M. in μ M (n = 3-6), or as a percentage of specific binding displaced at 100 μ M.
- c Displacement of specific [¹²⁵I]AB-MECA binding at human A₃ receptors expressed in HEK293 cells, in membranes, expressed as $K_i \pm$ S.E.M. in μ M (n = 2-5), or as a percentage of specific binding displaced at 100 μ M.
- d Displacement of ≤ 10 % of specific binding at the indicated concentration in M.
- e 2,3-trans
- f Chlorine was introduced in the 3-position using sulfonyl chloride in chloroform. Chem. Ind. London, 937 (1980)

EXAMPLE 17

This Example further illustrates the efficacy of the compounds of the present invention in blocking adenosine receptors. The affinities of the flavonoid derivatives 12-
5 16 were determined using radioligand binding assays, and the results thereof are set forth in Table 4.

Table 4. Affinities of flavonoid derivatives in radioligand binding assays at A_1 , A_{2a} , and A_3 receptors.^{a,c} $R_2=H$



Compound	R_3	R_1	R_4	rA_1^a	rA_{2a}^b	hA_3^c	rA_1/hA_3
12 flavanone	H	H	Ph	32.0±4.8	--	50.1±27.1	0.64
13	H	H	2'-OH-Ph	2.64±0.56	17.6±2.10	6.07±1.43	0.43
14 ^e	OH	H	2'-OH-Ph	91.6	d (10 ⁻⁴)	27±3% (10 ⁻⁴)	<1
15 ^e	OH	6-OMe	CH=CH-Ph	d (10 ⁻⁴)	d (10 ⁻⁴)	21.1±9.9	>8
16 ^e	OH	6-OMe	C≡C-Ph	50.3±17.0	d (10 ⁻⁴)	8.17±0.43	6.2


a Displacement of specific [³H]PIA binding in rat brain membranes, expressed as $K_i \pm$ S.E.M. in μ M ($n = 3-5$), or as a percentage of specific binding displaced at 100 μ M.

- b Displacement of specific [³H]CGS 21680 binding in rat striatal membranes, expressed as $K_i \pm$
S.E.M. in μM ($n = 3-6$), or as a percentage of specific binding displaced at 100 μM .
- c Displacement of specific [¹²⁵I]AB-MECA binding at human A_1 receptors expressed in HEK293
5 cells, in membranes, expressed as $K_i \pm$ S.E.M. in μM ($n = 2-5$), or as a percentage of
specific binding displaced at 100 μM .
- d Displacement of ≤ 10 % of specific binding at the indicated concentration in M.
- 10 e 2,3-trans

EXAMPLE 18

This Example further illustrates the efficacy of the compounds of the present invention in blocking adenosine receptors. The affinities of the flavonoid derivatives 17-
5 24 were determined using radioligand binding assays, and the results thereof are set forth in Table 5.

Table 5. Affinities of flavonoid derivatives in radioligand binding assays at A_1 , A_{2A} and A_3 receptors.^{a-d} $R_3=H$




Compound	R_1	R_4	rA_1^a	rA_{2A}^b	hA_3^c	rA_1/hA_3
17 visnagin	OCH_3	CH_3	$36 \pm 3\% (10^{-4})$	42.4 ± 11.9	60.0 ± 17.8	~ 2
18	OCH_3	CHO	$d (10^{-4})$	$25 \pm 8\% (10^{-4})$	88.9 ± 27.1	> 2
19	OCH_3	$CH=CH-Ph$	32.6 ± 10.5	11.5 ± 1.3	8.28 ± 2.69	0.37
20	OC_2H_5	$CH=CH-Ph$	35.6 ± 12.7	33.8 ± 14.8	1.16 ± 0.45	31
21	$O(CH_2)_2CH_3$	$CH=CH-Ph$	40.0 ± 3.5	49.0	3.95 ± 1.98	10
22	OCH_3	$CH=CH-CH=CH-Ph$	$d (10^{-4})$	$d (10^{-4})$	$d (10^{-4})$	---
23	OC_2H_5	$CH=CH-CH=CH-Ph$	$d (10^{-4})$	167	45.5 ± 10.3	> 4
24	OCH_3	$CH=NH-Ph$	$d (10^{-4})$	$d (10^{-4})$	9.18 ± 2.56	> 20

- a Displacement of specific [³H]PIA binding in rat brain membranes, expressed as $K_i \pm$ S.E.M. in μ M (n = 3-5), or as a percentage of specific binding displaced at 100 μ M.
- 5 b Displacement of specific [³H]CGS 21680 binding in rat striatal membranes, expressed as $K_i \pm$ S.E.M. in μ M (n = 3-6), or as a percentage of specific binding displaced at 100 μ M.
- 10 c Displacement of specific [¹²⁵I]AB-MECA binding at human A₂ receptors expressed in HEK293 cells, in membranes, expressed as $K_i \pm$ S.E.M. in μ M (n = 2-5), or as a percentage of specific binding displaced at 100 μ M.
- d Displacement of ≤ 10 % of specific binding at the indicated concentration in M.

EXAMPLE 19

This Example further illustrates the efficacy of the compounds of the present invention in blocking adenosine receptors. The affinities of the flavonoid derivatives 25-
5 39 were determined using radioligand binding assays, and the results thereof are set forth in Table 6.

Table 6. Affinities of flavone analogues determined in radioligand binding assays at A_1 , A_{2a} and A_3 receptors.

K_2 (μ M) or % inhibition at the indicated conc. (M)

Compound	R_1, R_2	R_3	R_4	$(rA_1)^a$	$(rA_{2a})^b$	$(hA_3)^c$
25	5-OH	H	Ph	2.17 \pm 0.06	6.20 \pm 1.24	47 \pm 12% (10^{-4})
26	7-OH	H	Ph	3.03 \pm 0.49	2.68 \pm 0.68	41 \pm 7% (10^{-4})
27	7-OH	H	3',4'-diMeO-Ph	18.8 \pm 3.6	35.4 \pm 7.3	32 \pm 4% (10^{-4})
28 apigenin	5,7-dioH	H	4'-OH-Ph	3.00 \pm 0.29	7.58 \pm 1.23	63 \pm 1% (10^{-5})
29	5-OH-7-Me	H	4'-OMe-Ph	3.40 \pm 0.35	28.0 \pm 7.35	6.70 \pm 1.78
30	5-OH-7-MeO	H	4'-OMe-Ph	d (10^{-4})	d (10^{-4})	69 \pm 7% (10^{-5})
31 tetramethyl- scutallarein	5,6,7-triMe	H	4'-Me-Ph	1.29 \pm 0.08	--	4.48 \pm 0.14
32 hispidulin	5,7-dioH-6-MeO	H	4'-OH-Ph	1.61 \pm 0.29	6.48 \pm 0.65	63 \pm 2%

Compound	R ₁ , R ₂	R ₃	R ₄	(rA ₁) ^a	(rA _{2a}) ^b	(hA ₃) ^c (10 ⁻⁵)
33 cirsimaritin	5-OH-6,7-diMeO	H	4'-OH-Ph	1.20±0.36	3.00±0.70	1.72±0.19
34	5,7-diAcO	OAc	H-Ph	11.6±4.4	56.5±9.5	17.5±2.0
35 tetramethyl- kaempferol	5,7-diMeO	OMe	4'-OMe-Ph	1.07±0.56	--	3.37±1.83
36 rhamnetin	7-OMe	OH	3',4'-diOH-Ph	--	--	1.38±0.18
37 quercetin	5,7-diOH	OH	3',4'-diOH-Ph	2.47±0.64	6.99±0.89	--
38 [*] pentamethyl morin	5,7-diMeO	OMe	2',4'-diOMe	27.6±7.5	46.7±2.7	2.65±0.72
39 hexamethyl myricetin	5,7-diMeO	OMe	3',4',5'-triOMe- Ph	d(10 ⁻⁶)	--	16.2±2.2

a Displacement of specific [³H]PIA binding in rat brain membranes, expressed as K_i ± S.E.M. in μM (n = 3-5).

5 b Displacement of specific [³H]CGS 21680 binding in rat striatal membranes, expressed as K_i ± S.E.M. in μM (n = 3-6).

- c Displacement of specific [125 I]AB-MECA binding at human A₁ receptors expressed in HEK-293 cells, in membranes, expressed as $K_i \pm$ S.E.M. in μ M ($n = 2-3$), or as a percentage of specific binding displaced at the specified conc. (M).
- 5 d Displacement of $< 10\%$ of specific binding at the specified conc. (M).
- e Also referred to as compound 6 in Table 3.

EXAMPLE 20

This Example further illustrates the efficacy of the compounds of the present invention in blocking adenosine receptors. The affinities of the flavonoid derivatives 40-51 were determined using radioligand binding assays, and the results thereof are set forth in Table 7.

Table 7. Affinities of compounds 40-51, determined in radioligand binding assays at A_1 , A_{2A} and A_3 receptors.^{a-d} For structures of the compounds, refer to Figures 8 and 9.

Compound	(rA_1) ^a	(rA_{2A}) ^b	(hA_3) ^c
40 genistein	5.0 ^e	36 ^e	20±2% (10 ⁻⁴)
41 (±) dihydro quercetin	d (10 ⁻⁵)	--	34.1±10.1
42 sakuranetin	--	--	3.40±0.18
43 α-naphtho- flavone	0.786± 0.018	1.32±0.43	71±2% (10 ⁻⁵)
44 β-naphtho- flavone	8.8±3.6	d (10 ⁻⁴)	12±1% (10 ⁻⁵)
45	7.10±0.10	29.9±10.1	d (10 ⁻⁴)
46	7.59±2.14	6.98±2.21	19±3% (10 ⁻⁴)
47 oxogalan- thinelactam	5.55±0.83	d (10 ⁻⁴)	d (10 ⁻⁴)
48 acetylhaem- anthamine methiodide	52.5±3.7	d (10 ⁻⁴)	d (10 ⁻⁴)
49	8.89±2.15	84.0±6.9	52±6% (10 ⁻⁴)
50 hema-toxylin	3.10±0.60	28±2% (10 ⁻⁴)	d (10 ⁻⁴)
51 arborinine	12.7±1.2	6.47±1.90	63±7% (10 ⁻⁴)

a Displacement of specific [³H]PIA binding in rat brain membranes, expressed as $K_i \pm$ S.E.M. in μ M (n = 3-5).

- b Displacement of specific [^3H]CGS 21680 binding in rat striatal membranes, expressed as $K_i \pm \text{S.E.M.}$ in μM ($n = 3-6$).
- 5 c Displacement of specific [^{125}I]AB-MECA binding at human A_1 receptors expressed in HEK-293 cells, in membranes, expressed as $K_i \pm \text{S.E.M.}$ in μM ($n = 2-3$), or as a percentage of specific binding displaced at the specified conc. (M).
- 10 d Displacement of $< 10\%$ of specific binding at the specified conc. (M).
- 15 e Okajima et al., Biochem. Biophys. Res. Comm., 203, 1488-1495 (1994)

EXAMPLE 21

- 20 This Example further illustrates the efficacy of the compounds of the present invention in blocking adenosine receptors. The effects of flavonoid derivatives 3a, 15, and 20 on the A_1 agonist-elicited inhibition of adenylyl cyclase were determined, and the results thereof are set forth in
- 25 Table 8.

Table 8. Effects of flavone derivatives 3a, 15, and 20 (50 μM) on the A_1 -agonist-elicited inhibition of adenylyl cyclase^a. Rat ($n=3$)

Compound	% Inhibition	
	IB-MECA (10^{-7}M)	IB-MECA (10^{-6}M)
galangin (control)	23.7 \pm 5.1	35.3 \pm 7.2
3a	2.2 \pm 2.2	19.1 \pm 7.4
15	25.9 \pm 2.4	39.2 \pm 3.0
20	21.2 \pm 3.9	40.9 \pm 3.7

- 35 a Assayed in membranes from CHO cells stably expressing the rat A_1 receptor in the presence of 1 μM forskolin.

EXAMPLE 22

This Example sets forth the sources for some of the flavonoids and dihydropyridines of the present invention.

Compounds 1, 11a, 12, and 17 were obtained from Fluka, Ronkonoma, NY or from Aldrich, St. Louis, MO. Compounds 13 and 14 were obtained from Apin Chemicals, Ltd., Oxon, UK. Compound 5a was obtained from K+K Laboratories, Jamaica, NY.

Compounds 5b (NSC #241010-z), 9 (NSC #78634-f), 11b (NSC #74876-t), 11c (NSC #74899-t), and 11d (NSC #74931-f) were obtained from NCI (Bethesda, MD).

Compounds 4, 27-34, 43, and 44 were obtained from Fluka, Ronkonoma, NY or from Aldrich, St. Louis, MO. Compound 36 was obtained from Apin Chemicals, Ltd., Oxon, UK. Compounds 25, 26, 27, and 42 were obtained from K+K Laboratories, Jamaica, NY. Compound 40 was obtained from Sigma, Milwaukee WI. Compound 47 was from Fisher, New York, NY. Compounds 46, 48 and 51 (1-hydroxy-2,3-dimethoxy-10-methyl-9(10H)-acridinone) were synthesized as reported in Literature. Fales et al.

J. Amer. Chem. Soc., 77, 5885-5890 (1955); Fales et al. Chem. & Ind., (London), 561-562 (1958); Craig et al., J. Org. Chem., 30, 1573-1576 (1965); Johne et al., Pharmazie, 25, 777-779 (1970); Banerjee et al., Bull. Calcutta Sch. Trop. Med., 13, 60 (1965).

R(+)- and S(-)-BAY K 8644, R(-)- and S(+)-niguldipine, nifedipine, nifedipine, nimodipine, and oxidized nifedipine were purchased from Research Biochemicals International (Natick, MA).

EXAMPLE 23

This Example illustrates a procedure for the oxidation (see Figure 7) of 1,4-dihydropyridine-3,5-dicarboxylate esters.

Equimolar amounts (0.25 mmol) of the 1,4-dihydropyridine-3,5-dicarboxylate ester and tetrachloro-1,4-benzoquinone in tetrahydrofuran (2 mL) were mixed and stirred for up to 4 h. The solvent was then evaporated, and products were purified by preparative TLC (silica 60; 1000

mm; Analtech, DE; 20% ethyl acetate - 80% petroleum ether 35-60).

EXAMPLE 24

5 This Example illustrates a procedure for the preparation (see Figure 6) of 1,4-dihydropyridine-3,5-dicarboxylate esters.

Equimolar amounts (0.5 mmol) of the appropriate 3-amino-2-butenate ester, aldehyde, and 3-ketopropionate ester derivative were dissolved in 5 mL of absolute ethanol. The solution was sealed in a glass tube and heated to 100 C for at least 24 h, and at most 72 h. The solvent was then evaporated, and products were purified either by crystallization or by preparative TLC (silica 60; 1000 mm; Analtech, DE; 20 % ethyl acetate - 80 % petroleum ether 35-60). From the moment the reactants were sealed into the glass tube, all procedures were performed under nitrogen and low-light conditions to prevent oxidation of the products. The products were shown to be pure by TLC.

20 Following the above procedure, several 1,4-dihydropyridine-3,5-dicarboxylate esters were prepared. The compounds thus prepared and their physical characterization data are set forth below. Additional characterization data are set forth in Table 9.

25

3-Methyl-5-ethyl 2,4,6-trimethyl-1,4(R,S)-dihydropyridine-3,5-dicarboxylate (52):

Yield: 98.2 % (recrystallized from MeOH); ¹H NMR (d-CHCl₃) δ 0.98 (d, 3H, 4-CH₃, J = 6.7 Hz); 1.31 (t, 3H, 5-methyl, J = 7.2 Hz); 2.28 (s, 6H, 2- & 6-CH₃); 3.73 (s, 3H, 3-methyl); 3.83 (q, 1H, H-4, J = 6.4 Hz); 4.13-4.26 (m, 2H, 5-methylene); 5.51 (wide signal, 1H, H-1). Molecular mass calcd.: 253.131 found: m/z = 271 (M + NH₄⁺, base), 254 (M + H⁺), CI.

**3-methyl-5-ethyl-
2,6-dimethyl-4-ethyl-1,4(R,S)-dihydropyridine-3,5-
dicarboxylate (56)**

- Yield: 50.5 %, recrystallized from MeOH. ¹H NMR
(d₆-DMSO) δ 0.63 (t, 3H, 4-methyl, J = 7.4 Hz); 1.18 (t, 3H,
5-methyl, J = 7.1 Hz); 2.19 (s, 6H, 2- & 6-CH₃); 3.58 (s, 3H,
3-methyl); 3.73 (t, 1H, H-4, J = 5.4 Hz); 4.00-4.13m, 4H, 4-
& 5-methylene). Molecular mass calcd.: 267.147 found: m/z =
285 (M + NH₄⁺, base), 268 (M + H⁺), CI.

**3-methyl,5-ethyl
2,6-dimethyl-4-propyl-1,4(R,S)-dihydropyridine-3,5-
dicarboxylate (57)**

- Yield: 73.3 %, recrystallized from MeOH. ¹H NMR
(d₆-DMSO) δ 0.77 (t, 3H, 4-methyl, J = 6.7 Hz); 1.12 (wide
signal, 4H, 4a- & 4b-methylene); 1.19 (t, 3H, 5-methyl, J =
7.2 Hz); 2.19 (s, 6H, 2- & 6-CH₃); 3.59 (s, 3H, 3-methyl);
3.76 (t, 1H, H-4, J = 5.2 Hz); 4.07 (q, 2H, 5-methylene, J =
6.6 Hz). Molecular mass calcd.: 281.163 found: m/z = 299 (M
+ NH₄⁺, base), 282 (M + H⁺), CI. Elemental analysis:
(C₁₅H₂₃N₁₀₄)C,H,N.

**3-methyl-5-ethyl-
2,6-dimethyl-4-(2(R,S),6-dimethyl-hexen-5-yl)-1,4(R,S)-
dihydropyridine-3,5-dicarboxylate (58)**

- Yield: 33.4 %, preparative TLC. ¹H NMR (d-CHCl₃) δ
0.91 (double d, 3H, 4-(2-CH₃), J₁ = 5.8 Hz, J₂ = 24.0 Hz);
1.30 (m, 7H); 1.63 (d, 6H, 4-(6- & 7-CH₃), J = 25.4 Hz); 2.28
(d, 6H, 2- & 6-CH₃, J = 7.8 Hz); 3.71 (s, 3H, 3-methyl);
3.97 (t, 1H, H-4, J = 7.2 Hz); 4.11-4.24 (m, 2H,
5-methylene); 5.07 (t, 1H, 4-(2-methyne), J = 6.5 Hz); 5.71
(wide signal, 1H, H-1). Molecular mass calcd.: 363.241
found: m/z = 381 (M + NH₄⁺, base), 364 (M + H⁺), CI.

**3-methyl,5-ethyl-
2,6-dimethyl-4-phenyl-1,4(R,S)-dihydropyridine-3,5-
dicarboxylate (59)**

- Yield: 78.6 %, recrystallized from MeOH. ¹H NMR
(d₆-DMSO) δ 1.11 (t, 3H, 5-methyl; J = 6.4 Hz); 2.23 (s, 6H,
2- & 6-CH₃); 3.51 (s, 3H, 3-methyl); 3.97 (m, 2H,
5-methylene); 4.84 (s, 1H, H-4); 7.08-7.20 (m, 5H,

4-phenyl). Molecular mass calcd.: 315.147 found: $m/z = 333$ ($M + NH_4^+$, base), 316 ($M + H^+$), CI.

5 **3-methyl-5-ethyl-2,6-dimethyl-4-(3-nitrophenyl)-1,4(R,S)-dihydropyridine-3,5-dicarboxylate (nitrendipine) (61)**

Yield: 88.3 %, preparative TLC. 1H NMR ($d-CHCl_3$) δ 1.25 (t, 3H, 5-methyl, $J = 5.4$ Hz); 2.39 (s, 6H, 2- & 6- CH_3); 3.66 (s, 3H, 3-methyl); 4.13 (m, 2H, 5-methylene); 5.11 (s, 1H, H-4); 5.71 (wide signal, 1H, H-1); 7.39 (t, 1H, H-5', $J = 8.0$ Hz); 7.65 (d, 1H, H-6', $J = 7.7$ Hz); 8.02 (d, 1H, H-4', $J = 7.8$ Hz); 8.13 (s, 1H, H-2'). Molecular mass calcd.: 360.132 found: $m/z = 378$ ($M + NH_4^+$, base), 361 ($M + H^+$), CI. Elemental analysis: $(C_{18}H_{20}N_2O_6)C, H, N$.

15 **3-methyl-5-ethyl-2,6-dimethyl-4-(4-nitrophenyl)-1,4(R,S)-dihydropyridine-3,5-dicarboxylate (67)**

Yield: 68.1 %, recrystallized from MeOH. 1H NMR (d_6 -DMSO) δ 1.11 (t, 3H, 5-methyl, $J = 7.2$ Hz); 2.26 (s, 6H, 2- & 6- CH_3); 3.52 (s, 3H, 3-methyl); 4.00 (m, 2H, 5-methylene); 4.96 (s, 1H, H-1); 7.39 (d, 2H, H-2' & H-6', $J = 8.7$ Hz); 8.09 (d, 2H, H-3' & H-5', $J = 8.7$ Hz). Molecular mass calcd.: 360.132 found: $m/z = 378$ ($M + NH_4^+$), base, 361 ($M + H^+$), CI.

25 **3-methyl-5-ethyl-2,6-dimethyl-4-(2- α,α,α -trifluoromethylphenyl)-1,4(R,S)-dihydropyridine-3,5-dicarboxylate (68)**

Yield: 45.6 %, preparative TLC. 1H NMR (d_6 -DMSO) δ 1.05 (m, 3H, 5-methyl); 2.21 (s, 6H, 2- & 6- CH_3); 3.44 (s, 3H, 3-methyl); 3.84-4.10 (m, 2H, 5-methylene); 5.38 (s, 1H, H-4); 7.31 (t, 1H, $J = 7.3$ Hz); 7.51 (m, 3H, 4-phenyl). Molecular mass calcd.: 383.134 found: $m/z = 401$ ($M + NH_4^+$, base), 384 ($M + H^+$), CI.

35 **3-methyl-5-ethyl-2,6-dimethyl-4-(4-methoxyphenyl)-1,4(R,S)-dihydropyridine-3,5-dicarboxylate (71)**

Yield: 61.8%, preparative TLC. 1H NMR ($d-CHCl_3$) δ 1.23 (t, 3H, 5-methyl, $J = 7.8$ Hz); 2.34 (s, 6H, 2- & 6- CH_3); 3.65 (s, 3H, 4'- OCH_3); 3.76 (s, 3H, 3-methyl); 4.07-4.14 (m, 2H, 5-methylene); 4.94 (s, 1H, H-4); 5.57 (wide signal, 1H,

H-1); 6.76 (d, 2H, H-2' & H-6', J = 8.5 Hz); 7.20 (d, 2H, H-3' & H-5', J = 8.6 Hz). Molecular mass calcd.: 345.158 found: m/z = 363 (M + NH₄⁺, base), 346 (M + H⁺), CI.

- 5 3-methyl-5-ethyl-2,6-dimethyl-4-(4-hydroxy,3-methoxyphenyl)-1,4(R,S)-dihydropyridine-3,5-dicarboxylate (72)

Yield: 79.6%, recrystallized from MeOH. ¹H NMR (d₆-DMSO) δ 1.13 (t, 3H, 5-methyl, J = 7.4 Hz); 2.21 (s, 6H, 2- & 6-CH₃); 3.52 (s, 3H, 3'-OCH₃), 3.66 (s, 3H, 3-methyl); 3.96-4.01 (m, 2H, 5-methylene); 4.74 (s, 1H, H-4); 6.49 (t, 10 1H, H-6', J = 3.9 Hz); 6.58 (d, 1H, H-5', J = 7.9 Hz); 6.66 (s, 1H, H-2'). Molecular mass calcd.: 361.153 found: m/z = 379 (M + NH₄⁺, base), 362 (M + H⁺), CI.

- 15 3-methyl-5-ethyl-2,6-dimethyl-4-[3,4-(methylenedioxy)phenyl]-1,4(R,S)-dihydropyridine-3,5-dicarboxylate (73)

Yield: 63.0%, preparative TLC. ¹H NMR (d-CHCl₃) δ 1.24 (t, 3H, 5-methyl, J = 7.3 Hz); 2.33 (s, 6H, 2- & 6-CH₃); 3.66 (s, 3H, 3-methyl); 3.73 (s, 1H, H-4); 4.09-4.22 (m, 2H, 20 5-methylene); 4.92 (s, 2H, 3',4'-methylenedioxy); 5.57 (wide signal, 1H, H-1); 5.89 (s, 1H, H-2'); 6.66 (d, 1H, H-5', J = 8.1 Hz); 6.75 (d, 1H, H-6', J = 7.9 Hz). Molecular mass calcd.: 359.137 found: m/z = 377 (M + NH₄⁺, base), 360 (M + H⁺), CI.

25

- 3-methyl-5-ethyl-2,6-dimethyl-4-[2-phenyl-(trans)-vinyl]-1,4(R,S)-dihydropyridine-3,5-dicarboxylate (75)

Yield: 87.9%, recrystallized from MeOH. ¹H NMR (d-CHCl₃) δ 1.31 (t, 3H, 5-methyl, J = 7.1 Hz); 2.34 (s, 6H, 30 2- & 6-CH₃); 3.74 (s, 3H, 3-methyl); 4.14-4.28 (m, 2H, 5-methylene); 4.63 (d, 1H, H-4, J = 5.4 Hz); 5.60 (wide signal, 1H, H-1); 6.19 (t, 1H, 4-(H-1 vinylidene), J = 6.0 Hz); 7.18 (d, 1H, 4-(H-2 vinylidene), J = 6.6 Hz); 7.24-7.34 (m, 5H, 4-phenyl). Molecular mass calcd.: 341.163 found: 35 m/z = 359 (M + NH₄⁺, base), 342 (M + H⁺), CI.

**3,5-diethyl-2,4-dimethyl-6-phenyl-1,4(R,S)-
dihydropyridine-3,5-dicarboxylate (78)**

Yield: 53%, recrystallized from petroleum ether (35-60). ¹H NMR (d-CHCl₃): δ 0.89 (t, 3H, 3-methyl, J = 7.4 Hz); 1.14 (d, 3H, 5-methyl, J = 7.4 Hz); 1.31 (t, 3H, J = 7.0 Hz); 2.3 (s, 3H, CH₃); 3.94 (m, 3H); 4.23 (m, 2H); 7.28-7.41 (m, 5H, C₆H₅). Molecular mass calcd.: 329.163 found: m/z = 347 (M + NH₄⁺, base), 330 (M + H⁺), CI.

**3,5-diethyl-2,4-dimethyl-6-butyl-
1,4(R,S)-dihydropyridine-3,5-dicarboxylate (77)**

The structure was confirmed by HNMR and mass spectrometry.

**3-methyl,5-(2-methoxy)ethyl
2,4,6-trimethyl-1,4(R,S)-dihydropyridine-3,5-dicarboxylate
(53)**

Yield: 53.0%; ¹H NMR (d-CHCl₃) δ 0.99 (d, 3H, 4-CH₃, J = 6.7 Hz); 2.28 (s, 6H, 2- & 6-CH₃); 3.41 (s, 3H, 5-methoxy); 3.66 (t, 2H, 5-(2-methylene), J = 4.9 Hz); 3.73 (s, 3H, 3-methyl); 3.85 (q, 1H, H-4, J = 6.5 Hz); 4.22-4.38 (m, 2H, 5-(1-methylene)); 5.54 (wide signal, 1H, H-1). Molecular mass calcd.: 283.142 found: m/z = 301 (M + NH₄⁺, base), 284 (M + H⁺), CI.

**3-methyl,5-benzyl 2,4,6-trimethyl-1,4(R,S)- dihydropyridine
-3,5-dicarboxylate (54)**

Yield: 70.5%; ¹H NMR (d-CHCl₃) δ 0.98 (d, 3H, 4-CH₃, J = 5.0 Hz); 2.28 (s, 6H, 2- & 6-CH₃); 3.72 (s, 3H, 3-methyl); 3.89 (q, 1H, H-4, J = 6.5 Hz); 5.21 (q, 2H, 5-methylene, J = 14.8 Hz); 5.54 (wide signal, 1H, H-1); 7.30-7.41 (m, 5H, 5-phenyl). Molecular mass calcd.: 315.147 found: m/z = 333 (M + NH₄⁺, base), 316 (M + H⁺), CI.

**3,5-Diethyl 2,6-dimethyl-4-(2-thienyl)-1,4-(±)-
dihydropyridine-3,5-dicarboxylate (114)**

¹H NMR (CDCl₃): δ 1.27 (t, 6H, J=7.75 Hz, 3 and 5-CH₂CH₃), 2.3 (s, 6H, 2 and 6-CH₃), 4.17 (m, 4H, 3 and 5-OCH₂), 5.35 (s, 1H, 4-H), 5.92 (br, 1H, NH), 6.8 (d, 1H, J=3.9 Hz, 3'-H), 6.85 (m, 1H, 4'-H), 7.06 (d, 1H, J=4.89 Hz, 5'-H).

3,5-Diethyl 2-methyl-4-phenylethynyl-6-trifluoromethyl-1,4-(±)-dihydropyridine-3,5-dicarboxylate

¹H NMR (CDCl₃): δ 1.32 (m, 6H, 3 & 5-CH₂CH₃); 2.35 (s, 3H, 7-CH₃), 4.20-4.38 (m, 4H, 3 & 5-OCH₂), 4.84 (s, 1H, 4-H); 6.29 (br, 1H, NH); 7.25-7.35 (m, 5H, C₆H₅). MS (EI: m/z 415 (M)⁺; 386 (M-C₂H₅)⁺; 342 (M-CO₂C₂H₅)⁺, base.

3,5-Diethyl 2-methyl-4,6-diphenyl-1,4-(±)-dihydropyridine-3,5-dicarboxylate (110).

¹H NMR (CDCl₃): δ 0.84 (t, 3H, J=6.83 Hz, 5-CH₂CH₃), 1.24 (t, 3H, J=6.84 Hz, 3-CH₂CH₃), 2.36 (s, 3H, 2-CH₃), 3.84 (m, 2H, 5-OCH₂), 4.12 (q, 2H, J= 6.83 Hz, 3-OCH₂), 5.12 (s, 1H, 4-H), 5.76 (br, 1H, NH), 7.18-7.44 (m, 10H, 4 and 6-C₆H₅).

3,5-Diethyl 2-methyl-4-(2-pyridyl)-6-phenyl-1,4-(±)-dihydropyridine-3,5-dicarboxylate (111).

¹H NMR (CDCl₃): δ 0.81 (t, 3H, J=6.84 Hz, 5-CH₂CH₃), 1.22 (t, 3H, J=6.83 Hz, 3-CH₂CH₃), 3.82 (m, 2H, 5-OCH₂), 4.11 (q, 2H, J=6.84 Hz, 3-OCH₂), 5.25 (s, 1H, 4-H), 5.88 (br, 1H, NH), 7.08-7.56 (m, 3H, pyridyl 3',4', and 5'-H), 7.37 (m, 5H, 6-C₆H₅), 8.55 (d, 1H, J=4.89, pyridyl 6'-H).

3,5-Diethyl 2-methyl-4-(3-pyridyl)-6-phenyl-1,4-(±)-dihydropyridine-3,5-dicarboxylate (112).

¹H NMR (CDCl₃): δ 0.84 (t, 3H, J=6.83 Hz, 5-CH₂CH₃), 1.24 (t, 3H, J=6.84 Hz, 3-CH₂CH₃), 2.38 (s, 3H, 2-CH₃), 3.83 (m, 2H, 5-OCH₂), 4.11 (q, 2H, J=6.84 Hz, 3-OCH₂), 5.09 (s, 1H, 4-H), 6.43 (BR, 1H, NH), 7.2-7.7 (m, 2H, pyridyl 4' and 5'-H), 7.3-7.4 (m, 5H, 6-C₆H₅), 8.33 (d, 1H, pyridyl 6'-H), 8.64 (s, 1H, pyridyl 2'-H).

**3,5-Diethyl 2-methyl-4-(4-pyridyl)-6-phenyl-1,4(±)-
dihydropyridine-3,5-dicarboxylate (113).**

¹H NMR (CDCl₃): δ 0.83 (t, 3H, J=6.84 Hz, 5-CH₂CH₃), 1.24 (t, 3H, J=6.84 Hz, 3-CH₂CH₃), 2.4 (s, 3H, -2-CH₃), 3.84 (m, 2H, 5-OCH₂), 4.12 (q, 2H, J=7.81 Hz, 3-OCH₂), 5.12 (s, 1H, 4-H), 6.35 (br, 1H, NH), 7.3-7.4 (m, 7H, 6-C₆H₅ and pyridyl 3' and 5'-H), 8.42 (d, 2H, J=5.86 Hz, pyridyl 2' and 6'-H).

**3,5-Diethyl 2-methyl-4-(2-benzofuryl)-6-phenyl-1,4-(±)-
dihydropyridine-3,5-dicarboxylate (116).**

¹H NMR (CDCl₃): δ 0.89 (t, 3H, J=6.84 Hz, 5-CH₂CH₃), 1.31 (t, 3H, J=6.84 Hz, 3-CH₂CH₃), 2.39 (s, 3H, -2-CH₃), 4.21 (m, 2H, 3-OCH₂), 5.48 (s, 1H, 4-H), 5.96 (br, 1H, NH), 6.48 (s, 1H, benzofuryl 3'-H), 7.1-7.5 (m, 9H, 6-C₆H₅ and benzofuryl 4', 5', 6' and 7'-H).

**3,5-Diethyl 2-methyl-4-(3-methyl phenylethynyl)-6-phenyl-
1,4-(±)-dihydropyridine-3,5-dicarboxylate (120).**

¹H NMR (CDCl₃): δ 0.95 (t, 3H, J=6.84 Hz, 5-CH₂CH₃), 1.35 (t, 3H, J=6.84 Hz, 3-CH₂CH₃), 2.29 (s, 3H, -3'-CH₃), 2.36 (s, 3H, 2-CH₃), 4.0 (m, 2H, 5-OCH₂), 4.3 (m, 2H, 3-OCH₂), 5.11 (s, 1H, 4-H), 5.92 (br, 1H, NH), 7.0-7.43 (m, 9H, 4-C₆H₄ and 6-C₆H₅).

25

**3-Ethyl 5-benzyl 2-ethyl-4-phenylethynyl-6-phenyl-1,4-
(±)-dihydropyridine-3,5-dicarboxylate**

¹H NMR (CDCl₃): δ 1.22 (t, J=6.9 Hz, 3H, 2-CH₂CH₃); 1.34 (t, J=6.9 Hz, 3H, 3-CH₂CH₃); 2.55, 3.01 (2m2H, 2-CH₂CH₃); 4.25 (m, 2H, 3-OCH₂); 5.06 (AB, J=12.7Hz, 2H, 5-OCH₂); 5.18 (s, 1H, 4-H); 5.95 (br, 1H, NH); 7.05-7.39 (m, 15H, 3 x C₆H₅)
MS (EI): m/z 491 (M)⁺; 462 (M-C₂H₅)⁺; 418 (M-CO₂C₂H₅)⁺; 356 (M-CO₂CH₂C₆H₅)⁺, base.

35

General Procedure for the Preparation of Compounds 122a-g, 125 and 126

Compound 139 (0.2 mmol) was dissolved in dry acetone
5 (10mL), anhydrous potassium carbonate (0.5 g) and
phenylethyl bromide or phenylpropyl bromide (5 eq.) were
added, and the mixture was refluxed for 2 h. The solution
were filtered and evaporated, and the residue were
separated with preparative TLC plates to give products
10 which were carried out for deprotection to afford 125 and
126 respectively.

**3-Ethyl 5-(2-methylbenzyl)-4-phenylethynyl-6-phenyl-
1,4(±)-dihydropyridine-3,5-dicarboxylate (122a)**

15 ¹H NMR (CDCl₃): δ 1.34 (t, 3H, J=6.84 Hz, 3-CH₂CH₃), 2.16 (s,
3H, 2'-CH₃), 2.37 (s, 3H, 2-CH₃), 4.26 (m, 2H, 3-OCH₂),
5.04 (AB, 2H, J=12.7 Hz, 5-OCH₂), 5.17 (s, 1H, 4-H), 5.92
(br, 1H, NH), 7.06-7.36 (m, 14H, 4-C₆H₅, 5-C₆H₄, and 6-
C₆H₅).

20

**3-Ethyl 5-(3-methylbenzyl)-4-phenylethynyl-6-phenyl-
1,4(±)-dihydropyridine-3,5-dicarboxylate (122b)**

¹H NMR (CDCl₃): δ 1.34 (t, 3H, J=6.84 Hz, 3-CH₂CH₃), 2.25 (s,
3H, 2'-CH₃), 2.36 (s, 3H, 2-CH₃), 4.3 (m, 2H, 3-OCH₂),
25 4.99 (AB, 2H, J=12.7 Hz, 5-OCH₂), 5.18 (s, 1H, 4-H), 5.91
(br, 1H, NH), 6.88-7.39 (m, 14H, 4-C₆H₅, 5-C₆H₄, and 6-
C₆H₅).

**3-Ethyl 5-(4-methylbenzyl)-4-phenylethynyl-6-phenyl-
1,4(±)-dihydropyridine-3,5-dicarboxylate (122c)**

30 ¹H NMR (CDCl₃): δ 1.34 (t, 3H, J=6.84 Hz, 3-CH₂CH₃), 2.37
(s, 3H, 2-CH₃), 4.3 (m, 2H, 3-OCH₂), 4.99 (AB, 2H, J=12.7
Hz, 5-OCH₂), 5.17 (s, 1H, 4-H), 5.86 (br, 1H, NH), 7.0-
7.4 (m, 14H, 4-C₆H₅, 5-C₆H₄, and 6-C₆H₅).

35

**3-Ethyl 5-(4-trifluoromethylbenzyl)-4-phenylethynyl-6-
phenyl-1,4(±)-dihydropyridine-3,5-dicarboxylate (122d)**

¹H NMR (CDCl₃): δ 1.35 (t, 3H, J=6.84 Hz, 3-CH₂CH₃), 2.37 (s, 3H, 2-CH₃), 4.3 (m, 2H, 3-OCH₂), 5.15 (AB, 2H, J=13.7 Hz, 5-OCH₂), 5.18 (s, 1H, 4-H), 5.94 (br, 1H, NH), 7.13-7.45 (m, 14H, 4-C₆H₅, 5-C₆H₄, and 6-C₆H₅).

5

3-Ethyl 5-(3-nitrobenzyl)-4-phenylethynyl-6-phenyl-1,4(±)-dihydropyridine-3,5-dicarboxylate (122f)

¹H NMR (CDCl₃): δ 1.35 (t, 3H, J=6.84 Hz, 3-CH₂CH₃), 2.38 (s, 3H, 2-CH₃), 4.3 (m, 2H, 3-OCH₂), 5.15 (AB, 2H, J=13.7 Hz, 5-OCH₂), 5.17 (s, 1H, 4-H), 5.94 (br, 1H, NH), 7.3-8.1 (m, 14H, 4-C₆H₅, 5-C₆H₄, and 6-C₆H₅).

10

3-Ethyl 5-(4-nitrobenzyl)-4-phenylethynyl-6-phenyl-1,4(±)-dihydropyridine-3,5-dicarboxylate (122g)

¹H NMR (CDCl₃): δ 1.36 (t, 3H, J=6.84 Hz, 3-CH₂CH₃), 2.38 (s, 3H, 2-CH₃), 4.3 (m, 2H, 3-OCH₂), 5.12 (AB, 2H, J=12.7 Hz, 5-OCH₂), 5.20 (s, 1H, 4-H), 5.94 (br, 1H, NH), 7.1-8.0 (m, 14H, 4-C₆H₅, 5-C₆H₄, and 6-C₆H₅).

15

20 3-Propyl 5-benzyl 2-methyl-4-phenylethynyl-6-phenyl-1,4(±)-dihydropyridine-3,5-dicarboxylate (124)

¹H NMR (CDCl₃): δ 1.01 (t, J=6.9 Hz, 3H, 3-CH₂CH₃), 1.73 (m, 2H, 3-CH₂CH₂); 2.37 (s, 3H, 2-CH₃); 4.15 (m, 2H, 3-OCH₂); 5.06 (AB, J=12.7 Hz, 2H, 5-OCH₂); 5.20 (s, 1H, 4-H); 5.88 (br, 1H, NH); 7.07-7.37 (m, 15H, 3 x C₆H₅) MS (EI: m/z 491 (M)⁺; 448 (M-C₃H₇)⁺; 404 (M-CO₂C₂H₅)⁺; 356 (M-CO₂CH₂C₆H₅)⁺, base.

25

30 3-Ethyl 5-phenylethyl 2-methyl-4-phenylethynyl-6-phenyl-1,4(±)-dihydropyridine-3,5-dicarboxylate (125).

¹H NMR (CDCl₃): δ 1.36 (t, J=6.8 Hz, 3H, 3-CH₂CH₃); 2.37 (s, 3H, 2-CH₃); 2.69 (t, J=6.8 Hz, 2H, 5-CH₂C₆H₅); 4.15 (m, 2H, 3-OCH₂); 4.30 (m, 2H, 5-OCH₂); 5.11 (s, 1H, 4-H); 5.87 (br, 1H, NH); 7.10-7.41 (m, 15H, 3 x C₆H₅) MS (EI: m/z 491 (M)⁺; 462 (M-C₂H₅)⁺; 418 (M-CO₂C₂H₅)⁺; 342 (M-CO₂(CH₂)₂(C₆H₅)⁺.

35

3-Ethyl 5-phenylpropyl 2-methyl-4-phenylethynyl-6-phenyl-1,4-(±)-dihydropyridine-3,5-dicarboxylate (126).

¹H NMR (CDCl₃): δ 1.30 (t, J=6.8 Hz, 3H, 3-CH₂CH₃); 1.65 (m, 2H, 5-CH₂CH₂CH₂); 2.35 (s, 3H, 2-CH₃); 2.40 (m, 2H, 5-CH₂C₆H₅); 3.98 (m, 2H, 3-OCH₂); 4.26 (m, 2H, 5-OCH₂); 5.14 (s, 1H, 4-H); 5.85 (br, 1H, NH); 7.20-7.41 (m, 15H, 3 x C₆H₅). MS (EI): m/z 505 (M)⁺; 476 (M-C₂H₅)⁺; 432 (M-CO₂C₂H₅)⁺; 342 (M-CO₂(CH₂)₃C₆H₅)⁺.

10

3-Ethyl 5-t-butyl 2-methyl-4-phenylethynyl-6-phenyl-1,4-(±)-dihydropyridine-3,5-dicarboxylate (127).

¹H NMR (CDCl₃): δ 1.25 (t, J=7.1 Hz, 3H, 3-CH₂CH₃); 1.31 (s, 9H, C(CH₃)₃); 2.31 (s, 3H, 2-CH₃); 4.22 (m, 2H, 3-OCH₂); 4.94 (s, 1H, 4-H); 5.56 (br, 1H, NH); 7.21-7.34 (m, 10H, 2 x C₆H₅). MS (EI): m/z 505 (M)⁺; 476 (M-C₂H₅)⁺; 432 (M-CO₂C₂H₅)⁺; 342 (M-CO₂(CH₂)₃C₆H₅)⁺.

15

3-(2-Methoxy-2-phenyl)-ethyl 5-Ethyl 2-methyl-4-phenylethynyl-6-phenyl-1,4-(±)-dihydropyridine-3,5-dicarboxylate (128).

20

¹H NMR (CDCl₃): δ 0.91 & 1.00 (2t, J=6.8 Hz, 3H, 5-CH₂CH₃); 2.35 (s, 3H, 2-CH₃); 4.00 (m, 2H, 5-OCH₂); 4.35 (m, 1H, 3-CH); 4.50 (m, 2H, 3-OCH₂); 5.05 (s, 1H, 4-H); 5.82 (br, 1H, NH); 7.21-7.41 (m, 10H, 2 x C₆H₅). MS (CI/NH₃): m/z 539 (M=NH₄)⁺.

25

3-Ethyl 5-(2-trimethylsilyl)-ethyl 2-methyl-4-phenylethynyl-6-phenyl-1,4-(±)-dihydropyridine-3,5-dicarboxylate (129).

¹H NMR (CDCl₃): δ 0.03 (s, 9H, Si(CH₃)₃); 0.79 (t, J=8.8 Hz, 2H, CH₂Si); 1.41 (t, J=6.8 Hz, 3H, 3-CH₂CH₃); 2.40 (s, 3H, 2-CH₃); 4.10 (t, J=8.8 Hz, 2H, 5-OCH₂); 4.31 (m, 2H, 3-OCH₂); 5.15 (s, 1H, 4-H); 5.87 (br, 1H, NH); 7.28-7.46 (m, 10H, 2 x C₆H₅). MS (CI/NH₃): m/z

35

3-Ethyl 5-thioethyl 2-methyl-4-phenylethynyl-6-phenyl-1,4(+)-dihydropyridine-3,5-dicarboxylate (130a)

Triethylamine (20 mg) was added to a mixture of compound 135 (90 mg, 0.2 mmol), diphenyl phosphoryl azide (56 mg, 0.2 mmol), and ethanethiol (20 mg, 0.3 mmol) in DMF (1 mL) with stirring and ice cooling. The mixture was stirred at room temperature for 3 hr, diluted with dichloromethane (20 mL), washed with water (10 mL x 2), dried with sodium sulfate. The solvent was evaporated and the residue was carried out for deprotection with 1N HCl to give 15 mg of product.

¹HMR (CDCl₃): 1.15(t, J=6.9 Hz, 3H, 5-CH₂CH₃); 1.28 (t, J=6.8 Hz, 3H, 3-CH₂CH₃); 2.37 (s, 3H, 2-CH₃); 2.82(m, 2H, 5-SCH₂); 4.31 (m, 2H, 3-OCH₂); 5.23 (s, 1H, 4-H); 5.95 (br, 1H, NH); 7.24-7.46(m, 10H, 2 x C₆H₅). MS (CI/NH₃); m/z

3-Ethoxycarbonyl-2-methyl-4-phenylethynyl-6-phenyl-1,4(+)-dihydropyridine-5-carboxylic acid (135)

¹H NMR (CDCl₃): 1.32(t, J=6.8 Hz, 3H, 3-CH₂CH₃); 2.38(s, 3H, 2-CH₃); 4.29(m, 2H, 3-OCH₂); 5.08(s, 1H, 4-H); 5.95(br, 1H, NH); 7.24-7.44(m, 10H, 2 x C₆H₅). MS (CI/NH₃); m/z 405 (M+NH₄); 388(MH)⁺.

3-Ethoxycarbonyl-2-methyl-4-phenylethynyl-6-phenyl-1,4(+)-dihydropyridine-5-carboxylic ethyl amide (132)

A mixture of compound 139 (75 mg, 0.17 mmol), N-hydroxysuccinimide (22 mg, 0.17 mmol) and DAEC (34 mg, 0.17 mmol) in DMF (1 mL) was stirred at room temperature for 4h. Ethylamine (2.0 M in THF, 0.4 mL) was added, and the reaction was stirred overnight. The solvent was removed and the residue was diluted with dichloromethane (10 mL), washed with water (5 mL x 2) and brine (5 mL x 2), dried with sodium sulfate. The solvent was evaporated and the residue was purified with preparative TLC plate to give 38 mg of compound 140 which was carried out for deprotection with 1N HCl to give 18 mg of product 132.

140: ^1H NMR (CDCl_3): 0.72 (t, $J=6.8$ Hz, 3H, 1- CH_2CH_3); 0.96 (t, $J=6.8$ Hz, 3H, 5- CH_2CH_3); 1.33 (t, $J=6.8$ Hz, 3H, 3- CH_2CH_3); 2.60 (s, 3H, 2- CH_3); 3.08 (m, 2H, 5- NCH_2); 3.62 (m, 2H, 1- OCH_2); 4.23 (m, 2H, 3- OCH_2); 4.45, 4.85 (AV, $J=11.7$ Hz, N- $\text{CH}_2\text{-O}$); 4.92 (s, 1H, 4-H); 4.97 (br, 1H, CONH); 7.22-7.44 (m, 10H, 2 x C_6H_5). MS (CI/ NH_3): m/z 473 (MH) $^+$.

132: ^1H NMR (CDCl_3): 0.85 (t, $J=6.8$ Hz, 3H 5- CH_2CH_3); 1.31 (t, $J=6.8$ Hz, 3H, 3- CH_2CH_3); 2.37 (s, 3H, 2- CH_3); 3.16 (m, 2H, 5- NCH_2); 4.25 (m, 2H, 3- OCH_2); 5.09 (s, 1H, 4-H); 5.37 (br, 1H, CONH); 5.64 (br, 1H, NH); 7.25-7.44 (m, 10H, 2 x C_6H_5). MS (CI/ NH_3): m/z

3-*t*-Butyl 5-ethyl 2-methyl-4-phenylethynyl-6-phenyl-1,4-
15 (+)-dihydropyridine-3,5-dicarboxylate (133)

^1H NMR (CDCl_3): δ 0.91 (t, $J=6.9$ Hz, 3H, 5- CH_2CH_3); 1.52 (s, 9H, $\text{C}(\text{CH}_3)_3$); 2.29 (s, 3H, 2- CH_3); 3.98 (m, 2H, 5- OCH_2); 5.05 (s, 1H, 4-H); 5.71 (br, 1H, NH); 7.21-7.40 (m, 10H, 2 x C_6H_5). MS (CI/ NH_3): m/z 461 ($\text{M} + \text{HN}_4$) $^+$; 443 (MH) $^+$.

20

5-Ethoxycarbonyl-2-methyl-4-phenylethynyl-6-phenyl-1,4-
(\pm)-dihydropyridine-3-carboxylic ethyl amide (134)

^1H NMR (CDCl_3): 0.85 (t, $J=6.8$ Hz, 3H, 3- CH_2CH_3); 1.18 (t, $J=6.8$ Hz, 3H, 5- CH_2CH_3); 2.24 (s, 3H, 2- CH_3); 3.38 (m, 2H, 3- NCH_2); 3.94 (m, 2H, 5- OCH_2); 4.80 (s, 1H, 4-H); 5.65 (br, 1H, NH); 6.32 (br, 1H, CONH); 7.23-7.38 (m, 10H, 2 x C_6H_5). MS (EI): m/z 414 (M) $^+$; 385 ($\text{M}-\text{C}_2\text{H}_5$) $^+$, base; 342 ($\text{M}-\text{CONHC}_2\text{H}_5$) $^+$.

3-Ethylcarbonyl 5-ethoxycarbonyl 2-methyl-4-
30 phenylethynyl-6-phenyl-1,4- \pm -dihydropyridine (135)

^1H NMR (CDCl_3): δ 0.93 (t, $J=7.0$ Hz, 3H, 5- CH_2CH_3); 2.33 (s, 3H, 2- CH_3); 2.48 (s, 3H, 3- COCH_3); 4.00 (q, 2H, $J=7.0$ Hz, 5- OCH_2); 5.04 (s, 1H, 4-H); 5.88 (br, 1H, NH); 7.26-7.43 (m, 10H, 2 x C_6H_5). MS (EI): m/z 385 (M) $^+$; 443 ($\text{M}-\text{CH}_3\text{CO}$) $^+$, base.

35

Ethyl 3-oxo-1,4,5,6,7,8-hexahydroquinoline-2-methyl-4-phenylethynyl-6-phenyl-1,4- \pm -dihydropyridin-5-carboxylate (136)

^1H NMR (CDCl_3); δ 1.02 (t, $J=6.8$ Hz, 3H, 5- CH_2CH_3); 2.04 (t, $J=4.9$ Hz, 2H, 7- CH_2); 2.46 (m, 4H, 8&9- CH_2); 4.00 (m, 2H, 5- OCH_2); 5.13 (s, 1H, 4-H); 6.14 (br, 1H, NH); 7.20-7.41 (m, 10H, 2 x C_6H_5). MS (EI: m/z 397 (M^+); 368 ($\text{M}-\text{C}_2\text{H}_5$) $^+$, base.

General procedure of N-H group protection in compound

10 (129):

Sodium hydride (60%) in mineral oil, 1.5 eq.) was added to compound 129 in solution of DMF (1.5 mL). The mixture was stirred for 5 min, chloromethyl methyl (ethyl) ether (1.5 eq.) was added slowly to the solution under argon at room temperature and stirred for 2h. The reaction was quenched by adding cold water (10 mL), extracted with ethyl acetate (10 mL x 2), the organic layer was washed with water (10 mL x 2), brine (10 mL x 2), dried with sodium sulfate. The solvent was evaporated and residue was purified with preparative TLC plates to give corresponding N-protected products 137 and 138.

1-Methoxymethyl 3-ethyl 5-(2-trimethylsilyl)-ethyl 2-methyl-4-phenylethynyl-6-phenyl-1,4-(\pm)-dihydropyridine-3,5-dicarboxylate (137)

^1H NMR (CDCl_3): δ 0.04 (s, 9H, $\text{Si}(\text{CH}_3)_3$); 0.63 (m, 2H, CH_2Si); 1.33 (t, $J=7.0$ Hz, 3H, 3- CH_2CH_3); 2.54 (s, 3H, 2- CH_3); 3.18 (s, 3H, OCH_3); 3.97 (t, $J=7.8$ Hz, 2H, 5- OCH_2); 4.27 (m, 2H, 3- OCH_2); 4.37, 4.85 (AB, $J=11.7$ Hz, N- $\text{CH}_2\text{-O}$); 5.07 (s, 1H, 4-H); 7.21-7.40 (M, 10H, 2 x C_6H_5). MS (CI/ NH_3): m/z 549 ($\text{M} + \text{NH}_4$) $^+$; 532 (MH) $^+$.

1-Ethoxymethyl 3-ethyl 5-(2-trimethylsilyl)-ethyl 2-methyl-4-phenylethynyl-6-phenyl-1,4-(\pm)-dihydropyridine-3,5-dicarboxylate (138)

¹H NMR (CDCl₃); δ-0.08 (s, 9H, Si(CH₃)₃); 0.61 (m, 2H, CH₂Si); 0.91 (t, J=6.9 Hz, 3H, OCH₂CH₃); 1.41 (t, J=6.9 Hz, 3H, 3-CH₂CH₃); 2.57 (s, 3H, 2-CH₃); 3.13 (m, 2H, OCH₂CH₃); 3.95 (t, J=7.9 Hz, 2H, 5-OCH₂); 4.12 (m, 2H, 3-OCH₂); 4.41, 4.81 (AB, J=10.8 Hz, 2H, N-CH₂-O); 5.02 (s, 1H, 4-H); 7.20-7.39 (m, 10H, 2 x C₆H₅). MS (CI/NH₃): m/z 563 (M+NH₄)⁺; 546 (MH)⁺.

1-Ethoxymethyl-3-ethoxycarbonyl-2-methyl-4-phenylethynyl-
10 6-phenyl-1,4-(±)-dihydropyridine-5-carboxylic acid (139)
TBAF (hydrate, 208 mg, 0.8 mmol) was added to a solution of 138 (115 mg, 0.21 mmol) in DMF (1 mL). The mixture was stirred under argon at room temperature for 2h, diluted with ethyl acetate (20 mL), washed with 1N HCl (5
15 mL), H₂O (20 mL x 2) and brine (20 mL x 2), dried with magnesium sulfate. The solvent was evaporated and residue was separated with preparative TLC plates to give 80 mg of product.

¹H NMR (CDCl₃); δ-0.93 (t, J=6.8 Hz, 3H OCH₂CH₃); 1.31 (t, J=6.9 Hz, 3H, 3-CH₂CH₃); 2.59 (s, 3H, 2-CH₃); 3.09, 3.65 (2m, 2H, OCH₂CH₃); 4.27 (m, J=7.9 Hz, 2H, 3-OCH₂); 4.39, 4.85 (2d, J= 10.7 Hz, 2H, 3-OCH₂); 4.41, 4.81 (AB, J=10.8 Hz, 2H, N-CH₂-O); 5.02 (s, 1H, 4-H); 7.12 (br, 1H, COOH); 7.20-7.39 (m, 10H, 2 x C₆H₅). MS (CI/NH₃): m/z 463 (M+NH₄)⁺; 446 (MH)⁺.

3,5-diethyl 2-methyl-4-(2-[4-nitro-phenyl]ethynyl)-
1,4(±)-dihydropyridine-3,5-dicarboxylate (119)

¹H NMR (d-CHCl₃) δ: 1.25-1.34 (m, 6H, 3-&5-CH₃); 2.49 (s, 3H, 2-CH₃); 3.47-3.59 (m, 2H, 3-CH₂); 4.23 (q, 2H, 5-CH₂, J=7.8 Hz); 5.42 (s, 1H, H-4); 56.29 (d, 1H, H-1, J=2.9 Hz); 7.55 (d, 2H, H-2' & H-6", J=8.8 Hz); 8.21 (d, 2H, H-3' & H-5', J=8.8 Hz); 8.26 (wide, 1H, H-6).

35 3,5-diethyl 2-methyl-4-(2-[3-toluyll]ethynyl)-1,4(±)-
dihydropyridine-3,5-dicarboxylate (120)

¹H NMR (d-CHCl₃) δ: 1.25-1.34 (m, 6H, 3-&5-CH₃); 2.49 (s, 3H, 2-CH₃); 3.47-3.59 (m, 2H, 3-CH₂); 4.23 (q, 2H, 5-CH₂, J=7.8 Hz); 5.42 (s, 1H, H-4); 56.29 (d, 1H, H-1, J=2.9 Hz); 7.55 (d, 2H, H-2' & H-6", J=8.8 Hz); 8.21 (d, 2H, H-3' & H-5', J=8.8 Hz); 8.26 (wide, 1H, H-6).

Table 8. Characterization data of dihydropyridine and derivatives.

Compound	T _m (°C)	formula	MS	elemental analysis	yield (%)	purification ^d
52	125-126	C ₁₃ H ₉ NO ₄	253 (CI)	C, H, N	98.2	1; MeOH
53	88-89	C ₁₄ H ₂₁ NO ₅ ·0.25H ₂ O	283 (CI)	C, H, N	53.0	2; TLC
54	oil	C ₁₈ H ₂₁ NO ₄ ·0.50H ₂ O	315 (CI)	C, H, N	70.5	2; TLC
56	83-84	C ₁₄ H ₂₁ NO ₄	267 (CI)	C, H, N	50.5	1; MeOH
57	95-96	C ₁₅ H ₂₃ NO ₄	281 (CI)	H, N ^a	73.3	1; MeOH
58	glass	C ₂₁ H ₃₃ NO ₄	363 (CI)	^b	33.4	2; TLC
59	121-122	C ₁₈ H ₂₁ NO ₄	315 (CI)	C, H, N	78.6	1; MeOH
61	78-79	C ₁₈ H ₂₀ N ₂ O ₆	360 (CI)	C, H, N	88.3	2; TLC
62	165-166	C ₁₉ H ₂₂ N ₂ O ₆	374 (EI)	C, H, N	35.5	1; EtOAc
67	151-152	C ₁₈ H ₂₀ N ₂ O ₆	360 (CI)	C, H, N	68.1	1; MeOH
68	114-115	C ₁₉ H ₂₀ F ₃ NO ₄	383 (CI)	C, H ^c	45.6	2; TLC

Compound	T _m (°C)	formula	MS	elemental analysis	yield (%)	purification ^d
71	112-113	C ₁₉ H ₂₁ NO ₅ -0.21 EtOAc	345 (CI)	C, H, N	61.8	2; TLC
72	190	C ₁₉ H ₂₃ NO ₆	361 (CI)	C, H, N	79.6	1; MeOH
73	oil	C ₁₉ H ₂₁ NO ₆ -0.32 EtOAc	359 (CI)	C, H, N	63.0	2; TLC
74	87-88	C ₂₀ H ₂₅ NO ₄	343 (CI)	C, H, N	71.9	2; TLC
75	135-136	C ₂₀ H ₂₃ NO ₄	341 (CI)	C, H, N	87.9	1; MeOH
76	176-177	C ₂₀ H ₂₁ NO ₄	339 (EI)	C, H, N	43.3	2; EtOH
77	oil	C ₁₇ H ₂₇ NO ₄	309 (CI)	C, H, N	58.8	2; TLC
78	123-124	C ₁₉ H ₂₃ NO ₄	329 (CI)	C, H, N	53.0	1; pe 35-60
79	oil	C ₂₆ H ₂₇ NO ₄	417 (CI)	C, H, N	34.7	2; column
80	oil	C ₁₃ H ₁₇ NO ₄	251 (EI)	C, H, N	57.9	2; TLC
81	oil	C ₁₃ H ₂₁ NO ₄ -0.20 EtOH	327 (EI)	C, H, N	48.9	2; TLC

a
b
5 c
d

57(C₁₅H₂₃NO₄) H,N;C: calcd, 64.03; found, 64.75;EI: calcd, 281.1627; found 281.1630
58(C₂₁H₃₃NO₄) C,H,N: calcd, 69.39,9.15,3.85; found, 61.65,7.27,4.70; EI: calcd,
363.2410; found
68(C₁₉H₂₉F₃NO₄) C,H,N: calcd, 3.65; found,4.17;EI: calcd,383.1344;found,383.1346
Purification was achieved either by:
1. recrystallization from the solvent specified, or

2. chromatography by the specified method, using EtOAc:pe 35-60 20:80((v/v)
eluent.


10

MeOH methanol
EtOH ethanol
EtOAc ethyl acetate
pe 35-60 petroleum ether 35-60 °C fraction
TLC preparative thin layer chromatography, silica 60, 1000 mm layer
thickness
column preparative column chromatography, silica 60, 220-440 mesh.


15

EXAMPLE 25

This Example illustrates the affinities of certain dihydropyridine derivatives. The affinities were
5 determined in radioligand binding assays, and the results thereof are set forth in Table 9.

Table 9. Affinities of dihydropyridine derivatives in radioligand binding assays at A_1 , A_{2a} , and A_3 receptors.

Compound	R ₂	R ₃	R ₄	R ₅	R ₆	rA ₁ ^a	rA _{2a} ^b	hA ₃ ^c
52	CH ₃	CO ₂ CH ₃	CH ₃	CO ₂ CH ₂ CH ₃	CH ₃	32.6±6.3	>100	32.3±5.1
53	CH ₃	CO ₂ CH ₃	CH ₃	CO ₂ (CH ₂) ₂ ⁻ -OCH ₃	CH ₃	49.2±0.7	19% (10 ⁻⁴)	62.3±16.7
54	CH ₃	CO ₂ CH ₃	CH ₃	CO ₂ CH ₂ Ph	CH ₃	6.45±1.47	9.72±0.63	2.78±0.89
55	CH ₃	CO ₂ CH ₂ CH ₃	CH ₃	CO ₂ (CH ₂) ₃ ⁻ -SPh	CH ₃	6.50±0.47	7.10±2.46	5.56±1.36
56	CH ₃	CO ₂ CH ₃	CH ₂ CH ₃	CO ₂ CH ₂ CH ₃	CH ₃	7.52±2.79	9.56±2.69	13.6±2.0
57	CH ₃	CO ₂ CH ₃	(CH ₂) ₂ CH ₃	CO ₂ CH ₂ CH ₃	CH ₃	8.17±1.58	11.5±3.8	6.51±0.74
58	CH ₃	CO ₂ CH ₃	CH ₂ CHCH ₃ (CH ₂) ₂ ⁻ CH=C(CH ₃) ₂ (R, S)	CO ₂ CH ₂ CH ₃	CH ₃	9.10±2.90	23.1±8.6	7.90±0.88

59	CH ₃	CO ₂ CH ₃	Ph	CO ₂ CH ₂ CH ₃	CH ₃	11.0±1.6	2.74±0.85	12.0±3.3
60	CH ₃	CO ₂ CH ₃	2-NO ₂ -Ph	CO ₂ CH ₃	CH ₃	2.89±0.23	18.2±2.51	8.29±2.41
nife- dipine								
61	CH ₃	CO ₂ CH ₃	3-NO ₂ -Ph	CO ₂ CH ₂ CH ₃	CH ₃	8.96±2.06	23.0±3.7	8.30±1.41
62	CH ₃	CO ₂ CH ₂ CH ₃	3-NO ₂ -Ph	CO ₂ CH ₂ CH ₃	CH ₃	3.34±2.17	18.2±79	2.51±0.15
63	CH ₃	CO ₂ CH ₃	3-NO ₂ -Ph	CO ₂ CH ₂ CH ₂ N -(Bz)CH ₃	CH ₃	19.6±1.9	63.8±4.2	3.25±0.26
64	CH ₃	CO ₂ CH(CH ₃) ₂	3-NO ₂ -Ph	CO ₂ CH ₂ CH ₂ O -CH ₃	CH ₃	20.1±1.7	44.3±14.4	8.47±2.75
65	CH ₃	CO ₂ CH ₃	3-NO ₂ -Ph	CO ₂ (CH ₂) ₃ -	CH ₃	41.3±3.5	~300	1.90±0.40
66	CH ₃	CO ₂ CH ₃	3-NO ₂ -Ph	 CO ₂ (CH ₂) ₃ -	CH ₃	d(10 ⁻⁴)	d(10 ⁻⁴)	2.80±0.35
67	CH ₃	CO ₂ CH ₃	4-NO ₂ -Ph	CO ₂ CH ₂ CH ₃	CH ₃	37±14% (10 ⁻⁴)	35.6±1.9	5.90±1.65
68	CH ₃	CO ₂ CH ₃	2-CF ₃ -Ph	CO ₂ CH ₂ CH ₃	CH ₃	6.68±2.37	20.7±2.8	11.6±1.7
69	CH ₃	CO ₂ CH ₃	2-CF ₃ -Ph	NO ₂	CH ₃	0.785± 0.113	35.1±10.1	2.77±0.34


70	CH ₃	CO ₂ CH ₃	2-CF ₃ -Ph	NO ₂	CH ₃	6.66±1.89	86.3±28.4	23.5±0.6
71	CH ₃	CO ₂ CH ₃	4-CH ₃ O-Ph	CO ₂ CH ₂ CH ₃	CH ₃	2.75±0.35	12.7±3.8	4.10±0.14
72	CH ₃	CO ₂ CH ₃	3-CH ₃ O-4-OH-Ph	CO ₂ CH ₂ CH ₃	CH ₃	51.0±3.7	56.8±1.9	32.1±9.2
73	CH ₃	CO ₂ CH ₃	3,4-OCH ₂ O-Ph	CO ₂ CH ₂ CH ₃	CH ₃	3.66±0.61	5.27±1.97	4.58±1.11
74	CH ₃	CO ₂ CH ₃	Ph-CH ₂ -CH ₂ -CH ₂	CO ₂ CH ₂ CH ₃	CH ₃	8.81±0.92	6.71±2.06	2.30±0.70
75	CH ₃	CO ₂ CH ₃	Ph-CH=CH-(trans)	CO ₂ CH ₂ CH ₃	CH ₃	16.1±0.5	49.3±12.5	0.670 ±0.195
76	CH ₃	CO ₂ CH ₃	Ph-C≡C-	CO ₂ CH ₂ CH ₃	CH ₃	5.39±0.33	38.3±7.9	0.940±0.070
77	CH ₃	CO ₂ CH ₂ CH ₃	CH ₃	CO ₂ CH ₂ CH ₃	(CH ₂) ₃ CH ₃	10.8±3.52	38.0±10.6	47.1±10.8
78	CH ₃	CO ₂ CH ₂ CH ₃	CH ₃	CO ₂ CH ₂ CH ₃	Ph	25.9±7.3	35.9±15.3	7.24±2.13
79	CH ₃	CO ₂ CH ₂ CH ₃	Ph-CH=CH-(trans)	CO ₂ CH ₂ CH ₃	Ph	5.93±0.27	4.77±0.29	0.108± 0.012

- a Displacement of specific [^3H]PIA binding in rat brain membranes, expressed as $K_i \pm \text{S.E.M.}$ in μM ($n = 3-5$).
- b Displacement of specific [^3H]CGS 21680 binding in rat striatal membranes, expressed as $K_i \pm \text{S.E.M.}$ in μM ($n = 3-6$).
- c Displacement of specific [^{125}I]AB-MECA binding at human A_1 receptors expressed in HEK cells, in membranes, expressed as $K_i \pm \text{S.E.M.}$ in μM ($n = 2-3$), or as a percentage of specific binding displaced at 10 μM .
- d Displacement of ≤ 10 % of specific binding at the indicated concentration in M.

EXAMPLE 26

This Example illustrates the affinities of certain pyridine derivatives. The affinities were determined in radioligand binding assays, and the results thereof are set forth in Table 10.

Table 10. Affinities of pyridine derivatives in radioligand binding assays at A_1 , A_{2a} , and A_3 receptors.



Compound	R ₂	R ₃	R ₄	R ₅	R ₆	rA ₁ ^a	rA _{2a} ^b	hA ₃ ^c
80	CH ₃	CO ₂ CH ₃	CH ₃	CO ₂ CH ₂ CH ₃	CH ₃	6.95±2.66	8.96±0.93	29.5±2.1
81	CH ₃	CO ₂ CH ₂ CH ₃	CH ₃	CO ₂ CH ₂ CH ₃	Ph	7.41±1.29	28.4±9.1	4.47±0.46
82	CH ₃	CO ₂ CH ₃	o-NO ₂ -Ph	CO ₂ CH ₃	CH ₃	d (10 ⁻⁴)	d (10 ⁻⁴)	d (10 ⁻⁴)
83	H	H	4-CH ₃ O-Ph	H	H	44.5±1.0	71±29	d (10 ⁻⁴)

a Displacement of specific [³H]PIA binding in rat brain membranes, expressed as $K_i \pm$ S.E.M. in μ M (n = 3-5).

b Displacement of specific [³H]CGS 21680 binding in rat striatal membranes, expressed as $K_i \pm$ S.E.M. in μ M (n = 3-6).

c Displacement of specific [¹²⁵I]AB-MECA binding at human A_3 receptors expressed in HEK cells, in membranes, expressed as $K_i \pm$ S.E.M. in μ M (n = 2-3), or as a percentage of specific binding displaced at 10 μ M.


d Displacement of ≤ 10 % of specific binding at the indicated concentration in M.

EXAMPLE 27

This Example illustrates the affinities of certain dihydropyridine and pyridine derivatives. The affinities were determined in radioligand binding assays, and the
5 results thereof are set forth in Tables 11-12.

Table 11. Affinities of certain dihydropyridine and pyridine derivatives in radioligand binding assays at A_1 , A_{2A} , and A_3 receptors.^{a-c}

5



86

K_i (μ M) or % inhibition^c

Compound	R_1	R_2	R_3	R_4	R_5	R_6	$\frac{IA_1}{IA_2}$	$\frac{IA_2}{IA_3}$	$\frac{IA_1}{IA_3}$
84	CH_2CH_3	$Ph-CH=CH-$ (trans)	CH_2CH_3	CH_2CH_3	CH_3	CH_3	4.65 ± 1.21	9.23 ± 3.60	0.887 ± 0.138
85	CH_2CH_3	$Ph-CH=CH-$ (trans)	CH_2CH_3	CH_2CH_3	CH_3	CH_3	13.7 ± 2.6	$14.4 \pm 4.8 (10^{-4})$	3.13 ± 0.51
86	CH_2CH_3	CH_2CH_3	CH_2CH_3	CH_2CH_3	Ph	Ph	21.5 ± 2.7	14.5 ± 3.5	8.49 ± 1.74
87	CH_2CH_3	CH_3	CH_2CH_3	CH_2CH_3	Ph	Ph	26.0 ± 8.7	3.15 ± 0.96	1.75 ± 0.47
88	CH_2CH_3	$Ph-CH=CH-$ (trans)	CH_2CH_3	CH_2CH_3	$4-CH_3-Ph$	$4-CH_3-Ph$	14.9 ± 4.9	$37 \pm 2 (10^{-4})$	9.13 ± 2.43
89	CH_2CH_3	$Ph-CH=CH-$ (trans)	CH_2CH_3	CH_2CH_3	$4-OCH_3-Ph$	$4-OCH_3-Ph$	9.49 ± 1.99	$d (10^{-4})$	1.43 ± 0.37
									5.2
									4.4
									2.5
									15
									1.6
									6.6

Compound	R_2	R_4	R_3	P_5	$\frac{rA_1}{hA_1}$	hA_3	$\frac{rA_1}{hA_1}$
90	CH_2CH_3	(trans) Ph-CH=CH- (trans)	CH_2CH_3	4-Cl-Ph	33.0 ± 7.5	$12 \pm 7\% (10^{-4})$	0.785 ± 0.272
91	CH_2CH_3	Ph-CH=CH- (trans)	CH_2CH_3	4-NO ₂ -Ph	13.1 ± 1.6		4.14 ± 0.51
92	CH_2CH_3	Ph-CH=CH- (trans)	CH_2CH_3	3-furyl	5.49 ± 0.25	$35 \pm 6\% (10^{-4})$	0.907 ± 0.307
93	CH_2CH_3	Ph-CH=CH- (trans)	CH_2CH_3	3-thienyl	7.52 ± 1.38	$d(10^{-4})$	0.407 ± 0.066
94	CH_2CH_3	2-OCH ₃ -Ph- CH=CH- (trans)	CH_2CH_3	Ph	25.3 ± 2.4	$17 \pm 12\% (10^{-4})$	0.334 ± 0.059
95	CH_2CH_3	2-NO ₂ -Ph-CH=CH- (trans)	CH_2CH_3	Ph	6.03 ± 1.39	$d(10^{-4})$	0.109 ± 0.017
96	CH_2CH_3	4-NO ₂ -Ph-CH=CH- (trans)	CH_2CH_3	Ph		0.07	
97	CH_2CH_3	(Ph) ₂ C=CH- Ph-C≡C-	CH_2CH_3	Ph	1.20 ± 0.14	8	1.42 ± 0.23
98	CH_2CH_3		CH_2CH_3	Ph	11.0 ± 0.1		0.0766 ± 0.015
99	CH_2CH_3	Ph-CH=CH- (trans)	CH_2Ph	Ph	$35 \pm 3\% (10^{-4})$	$15 \pm 3\% (10^{-4})$	0.0583 ± 0.012
100	CH_2CH_3	4-NO ₂ -Ph-CH=CH- (trans)	CH_2Ph	Ph	$33 \pm 1\% (10^{-4})$	$d(10^{-4})$	0.0724 ± 0.037
101	CH_2CH_3	Ph-C≡C-	CH_2Ph	Ph	40.1 ± 7.5	$d(10^{-4})$	0.0314 ± 0.002

Compound	R ₂	R ₁	R ₃	R ₄	R ₅	$\frac{RA_1}{A_1}$	$\frac{RA_{2A}}{A_{2A}}$	$\frac{hA_3}{8}$	$\frac{RA_1/hA_3}{1}$
102	CH ₂ CH ₃	4-NO ₂ -Ph-C≡C	CH ₂ Ph		Ph			<0.1	
103	CH ₂ CH ₃	4-NH ₂ -Ph-C≡C-	CH ₂ Ph		Ph			<0.1	
104	CH ₂ CH ₃	4-NH ₂ -3-I-Ph- C=C-	CH ₂ Ph		Ph			<0.1	
105	CH ₂ Ph	Ph-CH=CH- (trans)	CH ₂ CH ₃		Ph	d(10 ⁻⁴)	16±11% (10 ⁻⁴)	0.142±0.047	>700
106	CH ₂ Ph	4-NO ₂ -Ph-CH=CH- (trans)	CH ₂ CH ₃		Ph	d(10 ⁻⁴)	d(10 ⁻⁴)	0.286±0.038	>400
107	CH ₂ Ph	Ph-C≡C-	CH ₂ CH ₃		Ph	24±4% (10 ⁻⁴)	d(10 ⁻⁴)	0.169±0.026	>600
A		Ph-CH=CH- (trans)	CH ₂ CH ₃			2.52	2.6	6.3	0.4
B		Ph-C≡C-	CH ₂ Ph			7.01±1.98	43±2% (10 ⁻⁴)	2.75±0.78	2.6

88

- a Displacement of specific [³H]PIA binding in rat brain membranes, expressed as K_d±S.E.M. in μM(n=3-5), or as a percentage of specific binding displaced at the indicated concentration (M).
- b Displacement of specific [³H]CGS 21680 binding in rat striatal membranes, expressed as K_d±S.E.M. in μM(n=3-5), or as a percentage of specific binding displaced at the indicated concentration (M).
- 5 c Displacement of specific [¹²⁵I]AB-MECA binding at human A₂ receptors expressed in HEK cells, in membranes, expressed as K_d±S.E.M. in μM(n=3-4), or as a percentage of specific binding displaced at the indicated concentration (M).
- d Displacement of ≤ 10% of specific binding at the indicated concentration (M).
- e values from van Rhee et al.

Table 12. Affinities of dihydropyridine derivatives in radioligand binding assays at A_1 , A_{2A} , and A_3 receptors.^{a,c}

Compound	R_2	R_4	R_5	K_i (μ M) or $\% inhibition^c$			
				RA_1^a	RA_{2A}^b	hA_3^c	$\Sigma A_2/hA_3$
108	OCH ₂ CH ₃	2-thienyl-	OCH ₂ CH ₃	4.48 \pm 0.90	25 \pm 5% (10 ⁻⁴)	8.56 \pm 1.22	0.52
109	OCH ₂ CH ₃	Ph-C \equiv C-	OCH ₂ CH ₃	23.6 \pm 2.9	25 \pm 3% (10 ⁻⁴)	1.58 \pm 0.56	15.0
110	OCH ₂ CH ₃	Ph	OCH ₂ CH ₃	8.03 \pm 2.05	d (10 ⁻⁴)	1.71 \pm 0.36	4.7
111	OCH ₂ CH ₃	2-pyridyl-	OCH ₂ CH ₃	17.4 \pm 16.0	25 \pm 3% (10 ⁻⁴)	12.2 \pm 0.3	1.4
112	OCH ₂ CH ₃	3-pyridyl-	OCH ₂ CH ₃	26.6 \pm 1.2	40 \pm 2% (10 ⁻⁴)	19.2 \pm 2.6	1.8
113	OCH ₂ CH ₃	4-pyridyl-	OCH ₂ CH ₃	14.9 \pm 0.2	44.7 \pm 3.7	8.31 \pm 1.84	1.8
114	OCH ₂ CH ₃	2-thienyl-	OCH ₂ CH ₃	8.59 \pm 1.90	36 \pm 4% (10 ⁻⁴)	1.04 \pm 0.26	8.3
115	OCH ₂ CH ₃	2-furyl-	OCH ₂ CH ₃	3.65 \pm 0.45	d (10 ⁻⁴)	0.314 \pm 0.056	12

108-136

137-140

R_2 = Methyl, R_5 = Phenyl

Compound	R ₂	R ₁	R ₂	\overline{IA}_1^a	\overline{IA}_{2A}^b	\overline{IA}_3^c	$\overline{IA}_1/\overline{IA}_3$
116	OCH ₂ CH ₃	2-benzofuryl-	OCH ₂ CH ₃	3.65±0.45	d(10 ⁻⁴)	0.314±0.056	12
117	OCH ₂ CH=CH ₂	4-NO ₂ -Ph-CH=CH (trans)	OCH ₂ CH ₃	d(10 ⁻⁴)	15±2% (10 ⁻⁴)	0.296±0.063	>300
118	OCH ₂ CH ₃	4-NO ₂ -Ph-CH=CH (trans)	OCH ₂ CH ₃	31±3% (10 ⁻⁴)	26±6% (10 ⁻⁴)	0.198±0.047	>500
119	OCH ₂ CH ₃	4-NO ₂ -Ph-C≡C-	OCH ₂ CH ₃	34.5±6.8	24±4% (10 ⁻⁴)	2.58±0.66	13
120	OCH ₂ CH ₃	3-CH ₃ -Ph-C≡C-	OCH ₂ CH ₃	41±5% (10 ⁻⁴)	d(10 ⁻⁴)	0.220±.108	>400
121	OCH ₂ CH ₃	Ph-C≡C-	OCH ₂ Ph	25±4% (10 ⁻⁴)	d(10 ⁻⁴)	0.0695±0.0131	1400
122a	OCH ₂ CH ₃	Ph-C≡C-	OCH ₂ (2-CH ₃)Ph	16±1% (10 ⁻⁴)	13±1% (10 ⁻⁴)	0.112±0.015	1000
122b	OCH ₂ CH ₃	Ph-C≡C-	OCH ₂ (3-CH ₃)Ph	d(10 ⁻⁴)	16±2% (10 ⁻⁴)	0.0524±0.017	>2000
122c	OCH ₂ CH ₃	Ph-C≡C-	OCH ₂ (4-CH ₃)Ph	d(10 ⁻⁴)	17±3% (10 ⁻⁴)	0.110±0.033	>1000
122d	OCH ₂ CH ₃	Ph-C≡C-	OCH ₂ (4-CF ₃)Ph	32±3% (10 ⁻⁴)	15±1% (10 ⁻⁴)	0.0177±0.0015	>5000
122e	OCH ₂ CH ₃	Ph-C≡C-	OCH ₂ (3-I)Ph	14±1% (10 ⁻⁴)	19±7% (10 ⁻⁴)	0.0937±0.0333	>1000
122f	OCH ₂ CH ₃	Ph-C≡C-	OCH ₂ (3-NO ₂)Ph	28±2% (10 ⁻⁴)	d(10 ⁻⁴)	0.00858±0.00426	>11,000
122g	OCH ₂ CH ₃	Ph-C≡C-	OCH ₂ (4-NO ₂)Ph	29±2% (10 ⁻⁴)	d(10 ⁻⁴)	0.00269±0.00096	>37,000
123	OCH ₂ CH ₃	Ph-C≡C-	OCH ₂ Ph-4-CO-NH(CH ₂) ₃ NH ₂			0.01	
124	OCH ₂ CH ₂ CH ₃	Ph-C≡C-	OCH ₂ Ph	27±7% (10 ⁻⁴)	16±2% (10 ⁻⁴)	0.0682±0.0149	>1400
125	OCH ₂ CH ₃	Ph-C≡C-	O(CH ₂) ₂ Ph	d(10 ⁻⁴)	d(10 ⁻⁴)	0.146±0.012	>1000
126	OCH ₂ CH ₃	Ph-C≡C-	O(CH ₂) ₃ Ph	d(10 ⁻⁴)	d(10 ⁻⁴)	0.0757±0.0258	>1300
127	OCH ₂ CH ₃	Ph-C≡C-	OC(CH ₃) ₃	25.3±2.7	27±5% (10 ⁻⁴)	3.10±0.64	8.2
128	OCH ₂ CH ₃	Ph-C≡C-	OCH ₂ CH(OCH ₃)Ph(R)	8.58±1.34	33±6% (10 ⁻⁴)	1.72±0.45	5.0
129	OCH ₂ CH ₃	Ph-C≡C-	O(CH ₂) ₂ Si(CH ₃) ₃	18.8±5.4	18±5% (10 ⁻⁴)	0.0596±0.0199	310
130a	OCH ₂ CH ₃	Ph-C≡C-	SCH ₂ CH ₃	53.0±13.6	17±3% (10 ⁻⁴)	0.567±0.185	93

Compound	R ₁	R ₂	R ₃	R ₄	R ₅	$\frac{R_2}{R_1}$ ^a	$\frac{R_3}{R_2}$ ^b	$\frac{R_4}{R_3}$ ^c	$\frac{R_5}{R_4}$ ^d
130b	CH ₂ CH ₃	Ph-C≡C-	OCH ₂ CH ₃	OH	30	5.15±1.83	0.290±0.082	~100	<1
131	OCH ₂ CH ₃	Ph-C≡C-	NHCH ₂ CH ₃	OH	8.20±0.40	16±6% (10 ⁻⁴)	5.56±1.69	2.9	22
132	OCH ₂ CH ₃	Ph-C≡C-	OCH ₂ CH ₃	OH	16.2±5.6	20±2% (10 ⁻⁴)	1.04±0.22	2.7	5.6
133	OC(CH ₃) ₃	Ph-C≡C-	OCH ₂ CH ₃	OH	23.1±1.6	17±7% (10 ⁻⁴)	2.27±0.81	28	
134	NHCH ₂ CH ₃	Ph-C≡C-	OCH ₂ CH ₃	OH	65.6±15.1	22±4% (10 ⁻⁴)	0.443±0.086		
135	CH ₃	Ph-C≡C-	OCH ₂ CH ₃	OH	12.6±1.9	27±4% (10 ⁻⁴)			
136	R ₃ -R ₄ = (CH ₂) ₃	Ph-C≡C-	OCH ₂ CH ₃	OH	12.5±1.5	27±4% (10 ⁻⁴)			
137	CH ₂ OCH ₃	Ph-C≡C-	OCH ₂ CH ₃	OH	13±3% (10 ⁻⁴)	19±5% (10 ⁻⁴)	n.d.	>1	
138	CH ₂ OCH ₂ CH ₃	Ph-C≡C-	OCH ₂ CH ₃	OH	19.1±3.1	36±7% (10 ⁻⁴)	>100	>1	
139	CH ₂ OCH ₂ CH ₃	Ph-C≡C-	OCH ₂ CH ₃	OH	18.5±2.3	35±1% (10 ⁻⁴)	10.9±0.9	1.7	
140	CH ₂ OCH ₂ CH ₃	Ph-C≡C-	OCH ₂ CH ₃	OH	18.5±2.3	35±1% (10 ⁻⁴)	10.9±0.9	1.7	

a Displacement of specific [³H]R-PIA binding in rat brain membranes, expressed as K_i±S.E.M. in μM (n=3-5), or as a percentage of specific binding displaced at the indicated concentration (M).

b Displacement of specific [³H]CGS 21680 binding in rat striatal membranes, expressed as K_i±S.E.M. in μM (n=3-6), or as a percentage of specific binding displaced at the indicated concentration (M).

c Displacement of specific [¹²⁵I]AB-MECA binding at human A₃ receptors expressed in HEK cells, in membranes, expressed as K_i±S.E.M. in μM (n=3-4).

d Displacement of ≤ 10% of specific binding at the indicated concentration (M).

EXAMPLE 28

This Example illustrates a method of synthesis and resolution of diastereomers of certain 1,4-dihydropyridines. The chemical synthesis and resolution are outlined in
5 Figures 9A-9B. Characterization of the 1,4-dihydropyridines derivatives is set forth in Table 13.

In order to synthesize a diastereomeric pair for resolution, the Hantzsch reaction was carried out using a chiral β -ketoester derived from a protected glycerol moiety
10 (205, Fig.9A). This intermediate consisted of an acetoacetate ester of (S)-(+)-2,2-dimethyl-1,3-dioxolane-4-methanol), of which the S-enantiomer, 205a, is shown in Fig. 9A. This β -ketoester and the other components of the reaction to form the 4-phenylethynyl-6-phenyl-1,4-
15 dihydropyridine derivative. Thus, 205a, propiolaldehyde, 206, and a benzoylacetoacetate ester, 207, were dissolved in ethanol and refluxed overnight. Proton NMR in deuterated chloroform showed the isolated, gummy product, 208a, to be a 1:1 mixture of S, S and R, S- diastereomers. The signals
20 from each of the two ester α -methylene groups were well resolved in the spectrum and separated by 0.1 ppm. The signals from the 4-H of the diastereomers were separated by 0.03 ppm. Subsequent resolution of the enantiomers (below) and addition of a chiral shift reagent verified that these
25 peaks corresponded to pure diastereomers.

Crystallization of the racemic mixture 208a from methanol provided a pure diastereoisomer of this 4-phenylethynyl-6-phenyl-1,4-dihydropyridine derivative. This isomer was shown by X-ray crystallography to be of the S, S
30 configuration. The remaining isomer was obtained from the mother liquor as an oil and was shown by ^1H -NMR to be pure. The ^1H -NMR resonances of the two separate isomers with and without the lanthanide shift reagent Eu(FOD) [see JACS, 1980, 102, 5903] were catalogued and are shown in Table 13.
35 Each of the isolated isomers was shown to be a pure diastereomer and with the peaks of the other isomer not visible in the spectrum. Without EuFOD the 4-H and a methylene resonances were downfield in the R,S vs the S,S


isomer. EuFOD (4 mg/mL) caused a downfield shift of all of the resonances. In each case the resonances of the S,S isomer were shifted downfield to a greater degree in the presence of the NMR shift reagent. For example, the signals from the 4-H were shifted by EuFOD by >0.7 ppm in the S,S isomer and by only 0.3 ppm in the R,S isomer. Thus, in the presence of EuFOD this important resonance was separated by 0.4 ppm between the two diastereomers. The resonances of the two ester α -methylene groups, which were the best resolved in the spectrum measured in the absence of EuFOD, in the presence of EuFOD were separated by ~0.3 ppm.

To establish the absolute configuration at the C4 position of compound 210a an x-ray crystallographic structure was determined. Small needles of 210a were grown by vapor diffusion from methanol/water solution. X-Ray diffraction studies were performed with a Siemens P4 diffractometer (MoK α radiation). Crystals of 210a are monoclinic (P2₁) with lattice parameters $a = 11.059(2)$ Å, $b = 8.212(2)$ Å, $c = 15.629(3)$ Å, $\beta = 104.46(1)^\circ$; and $Z = 2$.

The carbonyl groups were in the cis/cis orientation, which is consistent with findings for other 1,4-dihydropyridines, in which the 4-position substituent is not highly hindered.

The 2,2-dimethyl-1,3-dioxolane moiety also served as a protected form of a diol, 211a obtained following deprotection in HCl/THF (Fig. 9B). This diol showed a selective reactivity vs the 5-ethyl ester in basic transesterification reactions. The 3-ethyl ester, 214, was obtained in this manner using sodium hydroxide in 95% ethanol. The resolved diastereomers 209a and 210a were also deprotected separately, to give 212a and 213a. These diols were then tested for biological activity. The results obtained are set forth in Table 14.

Table 13: Chemical shifts of diastereomeric 1,4-dihydropyridine derivatives in proton NMR.^a



209a(S,S), 210a(R, S)

5


Chemical shift (ppm from TMS)^a

Resonance	209a	209a + Eu(fod) ₃	210a	210a + Eu(fod) ₃
2-CH ₃	2.37	2.59	2.37	2.49
4-H	5.10	5.83	5.13	5.43
3-OCH ₂	4.26	4.79	4.36	4.46
5-OCH ₂	3.99	4.47	4.09	4.19
N-H	5.90	6.03	5.88	5.94

10

a in CDCl₃, in the presence or absence of 4 mg/mL Eu(fod)₃ (europium tris(6,6,7,7,8,8,8,-heptafluoro2,2-dimethyl-3,5-octanedionate)). The dihydropyridines were dissolved at a concentration of 5 mg/mL.

Table 14. Affinities of 1,4 dihydropyridine derivatives in radioligand binding assays at A_1 , A_{2A} , and A_3 receptors.^{a,c}





5

95

K_i (μ M) or $\% inhibition^d$

Compound	R_2	chirality at 4-position	R_3	\bar{rA}_1^a	\bar{rA}_{2A}^b	hA_3^c	\bar{rA}_1/hA_3
101	OCH_2CH_3	racemic	OCH_2Ph	40.1 ± 7.5	$d(10^{-4})$	0.0314 ± 0.0028	1300
209a		S	OCH_2CH_3	$32 \pm 4\%$ (10-4)	$d(10^{-4})$	1.31 ± 0.57	>70
210a		R	OCH_2CH_3	16.4 ± 3.1	$d(10^{-4})$	0.426 ± 0.133	38
212a		S	OCH_2CH_3	6.83 ± 0.20	$d(10^{-4})$	0.65	11

Compound	R ₂	chirality at 4-position	R ₃	$\frac{rA_1}{rA_2}$ ^a	$\frac{rA_{2b}}{rA_{2c}}$ ^b	$\frac{hA_3}{hA_1}$ ^c	$\frac{rA_1}{hA_1}$
213a		R	OCH ₂ CH ₃	14.8±1.0	39±8% (10 ⁻⁴)	0.52	28
	OCH ₂ CH ₃	racemic	OCH ₂ CH ₃	11.0±0.1	26±12% (10 ⁻⁴)	0.0766±0.0151	140
		racemic	OCH ₂ Ph				
	OCH ₂ CH ₃	racemic	OCH ₂ (4-CF ₃)Ph				
	OCH ₂ CH ₃	racemic	OCH ₂ CH(OCH ₃)-Ph (R)	8.58±1.34	33±6% (10 ⁻⁴)	1.72±0.45	5.0

a Displacement of specific [³H]R-PIA binding in rat brain membranes, expressed as K_d±S.E.M. in μM(n=3-5), or as a percentage of specific binding displaced at the indicated concentration (M).

b Displacement of specific [³H]CGS 21680 binding in rat striatal membranes, expressed as K_d±S.E.M. in μM(n=3-6), or as a percentage of specific binding displaced at the indicated concentration (M).

5 c Displacement of specific [¹²⁵I]AB-MECA binding at human A₃ receptors expressed in HEK cells, in membranes, expressed as K_d±S.E.M. in μM(n=3-4).

d Displacement of ≤ 10% of specific binding at the indicated concentration (M).

e values taken from van Rhee et al.¹⁶ or from Jiang et al.¹⁴

EXAMPLE 29

This Example illustrates the synthetic methodology adapted to prepare the 5-N-acyl derivatives of triazoloquinazolines. CGS15943 refers to compound 144 wherein R₁ and R₂ are hydrogen.

General Procedure for Preparation of 5-N-acyl derivatives of CGS15943

Method A (Symmetrical Anhydride)

To a stirred solution of CGS15943 (10 mg, 0.035 mmole), anhydride (0.105 mmole) and dimethylaminopyridine (0.5 mg, 0.004 mmole) in 1.5 ml of anhydrous DMF was added triethylamine (73 μ l, 0.525 mmole) at room temperature. The mixture was stirred for 48 h and then evaporated to dryness under reduced pressure. The residue was purified by preparative silica gel TLC (CH₂Cl₂-MeOH, 50:1 ~75:1) to afford the desired compounds, 145-148, and 150, 151.

Method B (Acid Chloride)

To a stirred solution of CGS15943 (10 mg, 0.035 mmole), anhydrous pyridine (40 μ l, 0.5 mmole) in 1.5 ml of anhydrous CH₂Cl₂ was added acylchloride (0.105 mmole) at 0°C. The mixture was stirred at room temperature for 24-48 h., then treated with same procedure as Method A for purification of desired compounds, (149, 153, and 160-162)

Method C (Carbodiimide)

A solution of CGS15943 (10 mg, 0.035 mmole), acid (0.210 mmole), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (41 mg, 0.201 mmole), 1-hydroxybenzotriazole (28 mg, 0.210 mmole), dimethylaminopyridine (0.5 mg, 0.004 mmole) and triethylamine (74 μ l, 0.530 mmole) in 2 ml of anhydrous DMF/CH₂Cl₂ (1:1 v/v) was stirred at room temperature for 48 h. The mixture was treated with same procedure as Method A for purification of desired compounds (152,154)

5-Acetylamino-9-chloro-2-(2-furyl)[1,2,4]triazolo[1,5-c]quinazoline (145).

2.78(3H, s), 6.63-6.65(1H, m), 7.30(1H, d, J=2.9), 7.68(1H, broad s), 7.73(1H, dd, J=1.9, 8.8), 7.86(1H, d, J=8.8), 8.48(1H, d, J=2.9), 8.99(NH, broad s)

9-Chloro-2-(2-furyl)-5-propionylamino[1,2,4]triazolo[1,5-c]quinazoline (146).

10

1.34(3H, t, J=7.81, 7.82), 3.10(2H, q, J=7.81, 6.8, 7.82), 6.63-6.65(1H, m), 7.31(1H, d, J=3.91), 7.68(1H, d, J=1.96), 7.73(1H, dd, J=1.96, 8.79), 7.88(1H, d, J=8.79), 8.48(1H, d, J=1.96), 9.01(NH, broad s)

15

5-Butyrylamino-9-chloro-2-(2-furyl)[1,2,4]triazolo[1,5-c]quinazoline (147)

1.10(3H, t, J=7.42, 7.28), 1.84-1.91(2H, m), 3.03(2H, t, J=7.42, 7.27), 6.63-6.65(1H, m), 7.31(1H, d, J=3.43), 7.68(1H, d, J=1.78), 7.73(1H, dd, J=2.34, 8.8), 7.88(1H, d, J=8.8), 8.48(1H, d, J=2.33), 8.97(NH, broad s)

9-Chloro-2-(2-furyl)-5-pentanoylamino[1,2,4]triazolo[1,5-c]quinazoline (148)

1.01(3H, t, J=6.84, 7.82), 1.48-1.55(2H, m), 1.77-1.85(2H, m), 3.05(2H, t, J=7.82, 6.83), 6.63-6.65(1H, m), 7.31(1H, d, J=2.93), 7.68(1H, broad s), 7.73(1H, dd, J=1.95, 8.79), 7.88(1H, d, J=8.80), 8.48(1H, d, J=2.93), 8.98(NH, broad s)

9-Chloro-2-(2-furyl)-5-trimethylacetylamino[1,2,4]triazolo[1,5-c]quinazoline (149).

1.47(9H, s), 6.63-6.65(1H, m), 7.31(1H, d, J=3.44), 7.68-7.69(1H, m), 7.74(1H, dd, J=2.47, 8.93), 8.01(1H, d, J=8.93), 8.49(1H, d, J=2.48), 9.39(NH, broad s)

35

5-[(tert.-Butoxycarbonyl)amino]-9-chloro-2-(2-furyl)[1,2,4]triazolo[1,5-c]quinazoline (150).

1.63 (9H, s), 6.65-6.66 (1H, m), 7.35 (1H, d, J=3.47), 7.69 (1H, broad s), 7.74 (1H, dd, J=2.60, 8.90), 8.01 (1H, d, J=8.90), 8.49 (1H, d, J=2.59), 8.56 (NH, broad s)

5 **5-Benzoylamino-9-chloro-2-(2-furyl)[1,2,4]triazolo[1,5-c]quinazoline (151).**

6.64-6.66 (1H, m), 7.35 (1H, d, J=2.93), 7.61-7.68 (3H, m), 7.68-7.70 (1H, m), 7.77 (1H, dd, J=1.96, 8.79), 8.04-8.10 (3H, m), 8.52 (1H, d, J=1.95), 9.75 (NH, broad s)

10

9-Chloro-2-(2-furyl)-5-(4'-iodobenzoyl)amino[1,2,4]triazolo[1,5-c]quinazoline (152).

6.64-6.66 (1H, m), 7.30-7.36 (3H, m), 7.70 (1H, broad s), 7.77 (1H, dd, J=1.95, 10.74), 7.92-8.10 (2H, broad m),

15 8.39 (1H, m), 8.52 (1H, d, J=1.96), 9.62 (NH, broad s)

9-Chloro-2-(2-furyl)-5-phenylacetyl amino[1,2,4]triazolo[1,5-c]quinazoline (153)

4.38 (2H, s), 6.62-6.65 (1H, m), 7.24-7.26 (1H, m), 7.35-7.44 (5H, m), 7.67-7.68 (1H, m), 7.74 (1H, dd, J=2.20, 9.04), 7.93 (1H, d, J=8.79), 8.49 (1H, m), 9.10 (NH, broad s)

20

5-[[4-tert.-Butoxycarbonyl]amino]butyryl]amino-9-chloro-2-(2-furyl)[1,2,4]triazolo[1,5-c]quinazoline (154).

25 1.44 (9H, s), 2.03 (2H, pen, J=6.83), 3.13 (2H, t, J=6.83, 7.33), 3.32 ((2H, q, J=6.35), 4.81 (NH, broad s), 6.63-6.65 (1H, m), 7.27-7.32 (1H, m), 7.67-7.68 (1H, m), 7.71-7.75 (1H, m), 7.91 (1H, d, J=8.79), 8.48 (1H, d, J=1.96), 9.15 (NH, broad s)

30

5-Amino-2-[2-5-bromofuryl]-9-chloro[1,2,4]triazolo[1,5-c]quinazoline (157).

A solution of 266 (0.01 g, 0.026 mmole) and N-bromosuccinimide (0.005 g, 0.028 mmole) in 2 ml of AcOH/CHCl₃,
35 (1:1) was stirred for 1 h at room temperature. The mixture was poured into 10 ml of sat. NaHCO₃ solution and the product was extracted with 10 ml of CHCl₃, three times. The combined

CHCl₃ solution was washed with brine, dried over anhydrous Na₂SO₄ and evaporated to dryness under reduced pressure. The residue was purified by preparative silical gel TLC (CHCl₃-MeOH, 80:1) to afford 2-[2-5-Bromofuryl]-5-[(tert.-butoxycarbonyl)amino]-9-chloro[1,2,4]triazolo[1,5-c]quinazoline (0.012 g, 99%) as a white solid : MS (CI, NH₃) 466(M⁺+1); ¹H-NMR (CDCl₃) 6.58 (1H, d, J=3.58), 7.29 (1H, d, J=3.59), 7.73 (1H, dd, J=2.28, 8.79), 7.98 (1H, d, J=9.01), 8.46 (1H, d, J=2.28), 8.53 (NH, broad s). To a solution of this compound in 2 ml of CH₂Cl₂ was added TFA (0.05 ml, 0.67 mmole) and stirred for 2 h at room temperature. The reaction mixture was treated with same work-up procedure above. A preparative silical gel TLC (n-Hex.-CHCl₃-MeOH, 1:1:0.1) of the crude product gave 273 (4.5 mg, 48%) as a white solid: MS (CI, NH₃) 366(M⁺+1), 383(M⁺+18); ¹H-NMR (CDCl₃) 5.94 (NH₂, broad s), 6.56 (1H, d, J=3.80), 7.25 (1H, d, J=3.80), 7.63-7.65 (2H, m), 8.41 (1H, d, J=2.07).

9-Chloro-2-(2-furyl)[1,2,4]triazolo[1,5-c]quinazolin-5-(6H)-one (158).

A solution of CGS15943 (0.075 g, 0.263 mmole) in 8.0 ml of AcOH and 2.0 ml of H₂O in a sealed tube was heated for 72 h at 100°C. The solution was coevaporated with toluene under reduced pressure and the residue was purified by preparative silical gel TLC (CHCl₃-MeOH, 15:1) to afford 274 (0.065 g, 86%) as a white solid : mp>310°C; MS (CI NH₃) 287(M⁺+1), 304(M⁺+18); ¹H-NMR (DMSO-d₆) 6.73-6.74 (1H, m), 7.26 (1H, d, J=3.47), 7.46 (1H, d, J=8.79), 7.77 (1H, dd, J=2.49, 8.90), 7.97 (1H, s), 8.16 (1H, d, J=2.48), 12.47 (NH, s).

9-Chloro-2-(2-furyl)[1,2,4]triazolo[1,5-c]quinazolin-6-propyl-5-one (159).


To a suspension of 158 (0.021 g, 0.073 mmole) in 2 ml of anhydrous THF was added a suspension of NaH (6 mg, 60% in mineral oil, prewashed with n-hexane, 0.15 mmole) in 2 ml of anhydrous THF followed by HMPA (0.32 ml, 12 mmole under N₂ atmosphere at room temperature. The mixture was stirred

vigorously for 30 min. H₂ gas evolved. 1-Bromopropane (28
μl, 0.3 mmole) was added and the reaction mixture was
refluxed for 6h. After cooling, the precipitate was removed
by filtration through a small volume of silical gel bed and
5 the filtrate was evaporated. The residue was purified by
preparative silical gel TLC (n-Hex.-EtOAc, 2:1) to afford
159 (0.012 g. 50%) as a white solid : ¹H-NMR (CDCl₃) 1.11(3H,
t, J=7.49, 7.48), 1.85-1.93(2H, m), 4.35(2H, t, J=8.03,
7.49), 6.60-6.61(1H, m); 7.32(1H, d, J=3.36), 7.37(1H, d,
10 J=9.34), 7.66(1H, broad s), 7.68(1H, dd, J=2.39, 9.01),
8.52(1H, d, J=2.50)

EXAMPLE 30

15 This Example illustrates the affinities of certain
triazoloquinazolines. The affinities were determined in
radioligand binding assays, and the results are set forth in
Tables 15-16.

Table 15: Affinities of 2-(2-furyl)[1,2,4]triazolo[1,5-c]quinazoline derivatives in radioligand binding assays at A_1 , A_{2A} , and A_3 receptors.^{a-c}



5

102

 K_i (μ M) or % inhibition^c

Compound	R_1	R_2	R_3	$\frac{rA_1}{hA_1}$	$\frac{rA_{2A}}{hA_{2A}}$	$\frac{rA_3}{hA_3}$	$\frac{rA_{2A}}{hA_3}$
144	H	H		21 \pm 3 ^d	3.3 \pm 1.7 ^d	13.8 \pm 2.4	
CGS15943							
145	COCH ₃	H		52.2 \pm 2.6		13.9 \pm 2.5	3.8
146	COCH ₂ CH ₃	H		283 \pm 42	106 \pm 30	7.66 \pm 3.03	40
147	CO(CH ₂) ₂ CH ₃	H		32.5 \pm 9.4	4	14.6 \pm 2.8	2.2
148	CO(CH ₂) ₃ CH ₃	H		28.9 \pm 3.7	16	21.5 \pm 6.2	1.3
149	COC(CH ₃) ₃	H		205 \pm 20	60	244 \pm 6	0.84
150	CO-OC(CH ₃) ₃	H		190 \pm 16	92.0 \pm 8.0	82.5 \pm 23.3	2.3
							1.1

Compound	R ₁	R ₂	R ₃	$\frac{rA_1}{rA_2}$	$\frac{hA_3}{rA_2}$	$\frac{rA_1}{hA_3}$	$\frac{rA_{2A}}{hA_3}$
151	CO-Ph	H		665±82	349±42	164	86
152	CO-(3-I-Ph)	H		200	200		
153	COCH ₂ -Ph	H		305±51	52.0±8.8		
154	CO(CH ₂) ₃ NH-Boc	H		91.3±13.2	173		
155	CO(CH ₂) ₃ NH ₂	H		11.3±2.4	14		
156	CO-Ph	Br		>1000	>1000		
157	H	Br		1,570 ^d	531 ^d		
158			H	3,000	4,000		
159			(CH ₂) ₂	2,000			
			CH ₃				


a Displacement of specific [³H]PIA binding in rat brain membranes, expressed as K_iS.E.M. in nM (n=3-5).

b Displacement of specific [³H]CGS 21680 binding in rat striatal membranes, expressed as K_iS.E.M. in nM (n=3-6).

c Displacement of specific [¹²⁵I]AB-MECA binding at human A₃ receptors expressed in HEK cells, in membranes, expressed as K_iS.E.M. in nM (n=3-4), or as a percentage of specific binding displaced at the indicated concentration (M).

5 d IC₅₀ values from Francis et al., J. Med. Chem., 31, pp. 1014-1020 (1988).

Table 16: Affinities of triazoloquinazoline derivatives in radioligand binding assays at A_1 , A_{2A} , and A_3 receptors.^{a-c}



5

104

K_i (μ M) or % inhibition^c

Compound	R_1	R_2	K_{A_1}	$K_{A_{2A}}$	hA_3	K_{A_1}/hA_3	$K_{A_{2A}}/hA_3$
160	CO-CH ₂ -CH ₂ -Ph	H	45.2 \pm 7.5	28.3 \pm 10.3	29.7 \pm 7.7		
161	CO-CH-CHPh	H	282 \pm 71	59.8 \pm 13.4	60.0 \pm 6.3		
162	CO-CH ₂ -Ph-4-NO ₂	H	28		8.70 \pm 0.97		
163	COCH ₂ -Ph-4-NH ₂	H	30	20	4		
164	COCH ₂ -Ph-3-I-4-NH ₂	H			1		
a	Displacement of specific [³ H]R-PIA binding in rat brain membranes, expressed as K_i \pm S.E.M. in nM (n=3-5).						
b	Displacement of specific [³ H]CGS 21680 binding in rat striatal membranes, expressed as K_i \pm S.E.M. in nM (n=3-6).						
c	Displacement of specific [¹²⁵ I]AB-MECA binding at human A_3 receptors expressed in HEK cells, in membranes, expressed as K_i \pm S.E.M. in nM (n=3-4), or as a percentage of specific binding displaced at the indicated concentration (M).						

10

EXAMPLE 31

This Example illustrates the utility of the adenosine receptor antagonists of the present invention in the killing of cancer cells.

5 Reagents:

HL-60 and U-937 cells were obtained from the ATCC (Bethesda, MD). RPMI 1640 medium and fetal bovine serum were obtained from Gibco BRL (Gaithersburg, MD). IB-MECA (2-[4-[92-carboxyethyl]phenyl]ethylamino]-5'-N-ethylcarboxamidoadenosine) and Cl-IB-MECA (N⁶-(3-iodobenzyl)-2-chloro-adenosine-5'-N-methyluronamide) were obtained from 10 RBI (Natick, MA) through the NIMH Synthesis Program. The A₁ selective adenosine antagonists 101 and 153 were synthesized using procedures described earlier. L-249313 (6-carboxymethyl-5,9-dihydro-9-methyl-2-phenyl-[1,2,4]triazolo[5,1-a][2,7]-naphthyridine) supplied as the mesylate salt was the gift of Dr. Marlene Jacobson at Merck (West Point, PA). The Apoptosis Detection System was from 15 Promega (Madison, WI), and the anti-bak antibody was from Calbiochem (La Jolla, CA). Rhodamine (TRITC)-conjugated AffiniPure Goat Anti-mouse IgG (H+L) is from Jackson ImmunoResearch Laboratories, Inc. Chromomycin A₃ was purchased from Sigma Chemical (St. Louis, MO).

25 Cell cultures and preparations:

The HL-60 cells were maintained in RPMI 1640 supplemented with 10% fetal calf serum, 100 units/ml penicillin, 100 mg/ml streptomycin and 2 mM L-glutamine. The cells were split every third day, and 2 days before each 30 experiment cultures were diluted to 2×10^5 cells/ml. For analysis of DNA content, aliquots of 2 ml were placed into 12-well flat-bottomed plates (Costar, Cambridge, MA, USA) containing 2-6 μ l test-compound solutions at defined concentrations or 6 μ l DMSO (diluting medium). Live cell 35 counting was carried out using 0.1% trypan blue.

DNA content analysis by flow cytometry:

Cells were fixed by adding $\sim 10^7$ cells suspended in 1 ml of PBS of 1 ml 80% ethanol at -20°C and stored for 48-120 hrs. After the cells were washed twice with PBS, the cells were stained with 20 mg/ml chromomycin A3 dissolved in PBS containing 2 mM MgCl_2 by incubation in subdued light (30 min; 4°C). The cells were then analyzed using a FACScan flow cytometer (Becton Dickinson, Mountain View, CA).

In situ hybridization:

Cells were fixed by immersing slides in 4% methanol-free formaldehyde in PBS at pH 7.4, 4°C , for 25 min. Fixed cells were permeabilized in 0.2% Triton[®] X-100 in PBS at 4°C for 5 min. After preincubation with 100 μM Promega equilibration buffer for 10 min at room temperature, the cells were covered with 50 μl TdT incubation buffer and a coverslip and maintained at 37°C for 60 min inside the dark humidified chamber to protect from dry and direct light. In order to stop the reaction, the coverslip was removed and the slides were immersed in 2xSSC for 15 min at room temperature. The cells were restained by propidium iodide 1 $\mu\text{g}/\text{ml}$ in PBS for 15 min. After every step, the cells were rinsed 2-3 times with fresh PBS for 5 min each. Finally, the glass slides were sealed with nail polish. The slides were stored at 4°C in the dark.

25

Immunofluorescent staining:

Cells were fixed as following steps: 10 min with 1% formalin in PBS, 5 min with 10°C methanol and 2 min with ice cold acetone, then permeabilized with 0.1% Nonidet P-40 for 20 min. After the preparation, the specimens were incubated step by step: 10% goat serum in PBS for 20 min, mouse monoclonal anti-bak antibody (5 $\mu\text{g}/\text{ml}$ PBS-BSA) for 60 min, Rhodamine (TRITC)-conjugated AffiniPure Goat Anti-mouse IgG (H+L) in PBS-BSA for 60 min. The specimens were washed 3-5 times with PBS after every step. The glass slides were stored at 4°C in the dark.

Flow cytometry data obtained are set forth in Figures 10A and 10B and indicate that high concentrations ($\geq \mu\text{M}$) of the A_1 receptor agonists, IB-MECA and Cl-IB-MECA, caused apoptosis in HL-60 promyelocytic leukemia cells. A similar response was observed in U-937 histiocytic lymphoma cells. In Figure 10, the percent of apoptotic cells was estimated from the percent cells having hypodiploid DNA content in a DNA frequency histogram. Upon incubation with the agonist for 48 hours, a plateau in the percent of apoptotic cells within the range of 30-50% of total was observed beginning at approximately 10 μM for either agonist in HL-60 (Figure 10A) and in U-937 cells (Figure 10B).

The more highly A_1 receptor selective agonist, Cl-IB-MECA, at lower concentrations ($\leq 1 \mu\text{M}$) did not significantly promote apoptosis, as determined using flow cytometry (Figures 10A-B), or general cell death, as determined using trypan blue staining (Figures 11A-D). In the latter assay, cell death was studied in HL-60 cells (Figures 11A-B) and in U-937 cells (Figures 11C-D) over a time course of 9 days. Cells in culture receiving no drug treatment proliferated geometrically until nearly the end of the experiment. In the presence of 0.01 or 1.0 μM Cl-IB-MECA alone, no deviation from this growth curve was observed.

Cl-IB-MECA at a concentration of 20 or 40 μM caused death of >90% of HL-60 cells in culture after 4 days followed by a very slow increase in cell number. U-937 cells were slightly less sensitive to this agonist, with 40 μM Cl-IB-MECA required to obtain a similar effect. Either of the selective A_1 receptor antagonists, 101 or L-249313, at a concentration of 0.5 μM , alone caused a general inhibition of proliferation of cells during this time period. An initial reduction in the number of cells during the first 4-5 days was followed by highly impeded growth. However, in the presence of both an A_1 antagonist and either 0.01 or 1.0 μM Cl-IB-MECA, the growth curves were nearly coincident with control curves. Thus, this A_1 agonist at sublethal concentrations, including the very low concentration of 10

nM, reversed the impairment of cell proliferation induced by either of the selective antagonists in both HL-60 cells (Figures 11A-B) and U-937 cells (Figures 11C-D).

The evidence that DNA strand breaks, characteristic of apoptosis, are induced by the A₂ antagonists was obtained using in situ end-labeling in-situ of fragmented nuclear DNA by terminal deoxynucleotidyl transferase (TUNEL method, Cooke, H., Trends in Biotech., 10, 23 (1992)), which incorporates a fluorescent label at the 3'-ends. In HL-60 and U-937 cells (Table 15), a 24 h incubation with either 50 nM 101 or 50 nM L-249313 produced widespread (approximately 30-40% of the cells present) fluorescent labeling of apoptotic cells. In the control cultures and in those treated with 10 nM Cl-IB-MECA alone, spontaneous apoptosis occurred in only 5-7% of the cells (Table 17).

Coadministration of 10nM Cl-IB-MECA with either antagonist suppressed the increase in apoptotic cells. The reversal of antagonist-induced apoptosis by the agonist is similar to that observed for the growth curves, except that both agonist and antagonist were applied at lower concentrations.

High concentrations of Cl-IB-MECA (30 μ M) also caused extensive DNA fragmentation in both HL-60 and U-937 cells, as indicated by the TUNEL method, with approximately 50% labeling.

The mechanism of the apoptosis observed in response to either A₂ agonists was investigated. Expression of the apoptosis-inducing protein bak in response to the adenosine receptor ligands was studied using immunofluorescent methods (Towbin et al., Proc. Natl. Acad. Sci. USA, 76, 4350-4354 (1979)). Bak labeling was absent in control HL-60 cells or U-937 cells, but was present in the cells treated for 24 hours with either L-249313 or 101. The expression of bak protein was also induced in both HL-60 cells and U-937 cells using the triazoloquinazoline A₂ antagonist 153 (10 nM). Cl-IB-MECA, at 30 μ M but not concentrations \leq 1 μ M, induced a significant level of expression of bak in both HL-60 cells and U-937 cells.

Various other cell lines were investigated for the expression of bak in response to the A₁ agonist. Cl-IB-MECA at 10 μ M was found to induce the expression of bak in MCF7 breast adenocarcinoma cells and 1321N astrocytoma cells, but
5 not in U373 astrocytoma cells, so the upregulation of bak expression by A₁ agonists appears to be a widespread but not universal phenomenon.

These results suggest that both A₁ agonists and antagonists, by virtue of regulating programmed cell death,
10 may have usefulness in treating diseases either in which cytotoxicity is undesirable, such as neurodegeneration, or desirable, such as cancer and inflammation. The level of agonist or antagonist should be carefully balanced to obtain the desired effect on the cells, e.g., death or protection.

Table 17: Percentage of apoptotic cell in HL60 and U937 after 48 hour treatment with A₁ antagonists (50 nM) and a low concentration of Cl-IB-MECA (10 nM), determined using end-labeling in-situ of fragmented nuclear DNA by terminal deoxynucleotidyl transferase. Both small and large cells that stained positive were counted manually.

Cell	Control	Cl-IB MECA	101	101 +Cl-IB-MECA	L-249313	L-249313 +Cl-IB-MECA
HL-60	6.8	7.1	39.7	11.6	31.2	12.0
U937	5.1	5.6	27.3	10.4	30.8	10.9

EXAMPLE 32

This Example illustrates the utility of the present inventive compounds in preserving neurons in stroke suffered by an animal.

5 Female Mongolian gerbils received bilateral 10 min. carotid occlusion, followed by injection 15 min. later with 100 µg/kg of the A₁ adenosine receptor antagonist, compound 101, in a vehicle of emulpher/saline (1:4). Control gerbils were injected with saline. The survival of the animals was
10 followed as a function of time after ischemia. The animals injected with the antagonist showed high survival rates: 90% after 7 days and 80% after 90 days. The control group survival rates were 50% after 7 days and 20% after 90 days.

 The number of intact neurons was also counted. After 7
15 days of ischemia, the animals injected with the antagonist had 75% intact neurons, and, after 90 days, the number of intact neurons was 90%, thereby indicating that neurons that are damaged under the influence of the antagonist are capable of recovery. The control group rates were 35% after
20 7 days and 90 days.

 These data show that the A₁ adenosine receptor antagonist both protects and enhances the chances of survival among neurons that are damaged and, unless exposed to the drug, would eventually die.

25

EXAMPLE 33

 This Example illustrates the bronchoconstricting effect of A₁ adenosine receptor agonists and the utility of A₁ adenosine receptor antagonists in combating the
30 bronchoconstricting effect of A₁ adenosine receptor agonists.

 Adenosine receptors have been implicated in the bronchoconstriction (BC) in allergic asthma. It was found that after selectively blocking A₁ receptors by diphenyl cyclopentyl xanthine, IB-MECA, an A₁ agonist, induced a
35 dose-dependent BC in allergic rabbits. It was also found that after blocking both A₁ and A₂ specific binding sites with a xanthine amino congener, A₁ specific binding was

displaced by APNEA, an A_1 agonist. These results confirmed that A_1 adenosine receptor agonist can cause BC.

It was further found that A_1 adenosine receptor antagonists can combat the BC induced by the A_1 agonist.

5 New Zealand white pasteurilla free rabbit litter mates were exposed to A_1 adenosine agonist APNEA, IB-MECA, or Cl-IB-MECA and the percent change in dynamic compliance (C_{dyn}) was measured as a function of the adenosine dose using known procedures. Ali et al., J. Pharmacol. & Exper. Therap.,
10 268, 1328-1334 (1993). Allergy was induced in the rabbits using dust mites.


The results obtained are set forth in Fig. 12 (open symbols) and show that the adenosine agonist causes the C_{dyn} to decrease, which is an indication of bronchoconstriction.
15 However, when an adenosine antagonist, compound 101, was co-administered with the agonist, the percent change in dynamic compliance was much reduced (filled symbols). The antagonist was administered at a constant dose of 10^{-5} M through a 2 minute aerosolization, followed by a 15 minute
20 wait before the measurement of dynamic compliance. The results confirm that the A_1 adenosine receptor antagonist is effective in combating inflammatory disorders such as asthma.

25 All of the references cited herein including patents, patent applications, and publications, are hereby incorporated in their entireties by reference.

While this invention has been described with an emphasis upon preferred embodiments, it will be obvious to
30 those of ordinary skill in the art that variations of the preferred embodiments may be used and that it is intended that the invention may be practiced otherwise than as specifically described herein. Accordingly, this invention includes all modifications encompassed within the spirit and
35 scope of the invention as defined by the following claims.

WHAT IS CLAIMED IS:

1. A compound of the formula



- 5 or a pharmaceutically acceptable salt thereof, wherein R_2 is a C_1 - C_6 alkyl; R_6 is selected from the group consisting C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, and phenyl which may be further substituted with C_1 - C_6 alkyl, halo, nitro, furyl, or thienyl; R_3 is selected from the group consisting of C_1 - C_6 alkyl, C_1 - C_6 alkyloxycarbonyl, aryl C_1 - C_6 alkyloxycarbonyl, C_1 - C_6 alkylthiocarbonyl, C_1 - C_6 alkylaminocarbonyl, and C_1 - C_6 alkyloxy C_1 - C_6 alkylcarbonyl, or R_3 together with R_2 forms a ring having 2-4 methylene groups, and C_1 - C_6 alkenyloxycarbonyl; R_4 is selected from 10 the group consisting of C_1 - C_6 alkyl, aryl C_2 - C_6 alkenyl, C_1 - C_6 alkylamino, C_1 - C_6 alkyl silyl C_1 - C_6 alkyloxy, aryl, heterocyclic, aryl C_1 - C_6 alkyl, phenylacetylenyl which may be further substituted with nitro, C_1 - C_6 alkyl, hydroxy, halo, amino, carboxy, C_1 - C_6 alkoxy, C_1 - C_6 haloalkyl, or C_1 - C_6 20 alkylamino, and styryl whose phenyl ring may be further substituted with one or more substituents selected from the group consisting of halo, nitro, amino, hydroxy, C_1 - C_6 alkyl, cyano, C_1 - C_6 alkyloxy, C_1 - C_6 alkyloxycarbonyl, C_1 - C_6 alkylcarbonyl, hydroxy C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, 25 carboxy, aminocarbonyl, C_1 - C_6 alkylamino, amino C_1 - C_6 alkyl, and C_1 - C_6 dialkylamino; and R_5 is selected from the group consisting of C_1 - C_6 alkyloxycarbonyl, aryl C_1 - C_6 alkyloxycarbonyl, C_1 - C_6 alkyloxy C_1 - C_6 alkyloxycarbonyl, aryloxy C_1 - C_6 alkyloxycarbonyl, C_1 - C_6 alkyloxycarbonyl, aryl 30 C_1 - C_6 alkyloxy C_1 - C_6 alkyloxycarbonyl, silyl C_1 - C_6 alkyloxycarbonyl, C_1 - C_6 alkylthio, hydroxy, and C_1 - C_6 alkylamino, wherein the aryl moiety of said R_5 may be

further substituted with C₁-C₆ alkyl, C₁-C₆ halo alkyl, trifluoromethyl, halo, nitro, C₁-C₆ amino alkyl, C₁-C₆ aminoalkylamino, or C₁-C₆ amino alkylamino carbonyl; wherein the aryl moiety of said R₃, R₄, R₅, and R₆ is
5 independently phenyl or naphthyl.

2. The compound of claim 1, wherein said heterocyclic is selected from the group consisting of furyl, thienyl, pyridyl, and benzofuryl.
10

3. The compound of claim 2, wherein R₂ is methyl.

4. The compound of claim 3, wherein R₃ is selected from the group consisting of methoxycarbonyl and
15 ethoxycarbonyl.

5. The compound of claim 4, wherein R₄ is selected from the group consisting of C₁-C₄ alkyl and phenyl.

20 6. The compound of claim 5, wherein R₄ is selected from the group consisting of C₁-C₃ alkyl.

7. The compound of claim 6, wherein R₅ is selected from the group consisting of methoxycarbonyl, ethoxycarbonyl, methoxyethoxycarbonyl, and
25 benzyloxycarbonyl.

8. The compound of claim 4, wherein R₄ is phenyl substituted with one or more substituents selected from the
30 group consisting of nitro, trifluoromethyl, methoxy, hydroxy, and methylenedioxy.

9. The compound of claim 8, wherein R₅ is selected from the group consisting of methoxycarbonyl, ethoxycarbonyl, and methoxyethoxycarbonyl.
35

10. The compound of claim 3, wherein R₃ is ethoxy, R₄ is phenylacetylenyl or trans-styryl, or phenylacetylenyl or

trans-styryl which is substituted with methoxy or nitro groups, R_5 is ethoxy or benzyloxy, and R_6 is methyl, ethyl, phenyl, benzyl, furyl, or thienyl wherein said phenyl, benzyl, furyl, and thienyl may be further substituted with
 5 nitro, halo, methyl, or methoxy group.

11. The compound of claim 10, wherein R_3 is ethoxy, R_4 is phenylacetylenyl, R_5 is benzyloxy, and R_6 is phenyl.


10 12. The compound of claim 1, wherein said compound is a mixture of diastereomers having R or S configuration at the 4-position.

13. The compound of claim 1, wherein said compound is
 15 an R-enantiomer.

14. The compound of claim 1, wherein said compound is an S-enantiomer.

20 15. The compound of any of claims 11-13, wherein said compound is modified as an acetoacetate ester of a (+) or (-) 2,2-dialkyl 1,3-dioxolane-4-methanol or a (+) or (-) alkylene glycol methanol.

25 16. A compound of the formula




or pharmaceutically acceptable salts thereof, wherein R_2 is selected from the group consisting of hydrogen and C_1 - C_6 alkyl; R_3 is selected from the group consisting of hydrogen and C_1 - C_6 alkyloxycarbonyl; R_4 is selected from the group
 30 consisting of C_1 - C_6 alkyl, phenyl C_2 - C_6 alkenyl, phenyl C_2 - C_6 alkynyl, aryl, and aryl substituted with one or more substituents selected from the group consisting of nitro and C_1 - C_6 alkyloxy; R_5 is selected from the group consisting of
 35 hydrogen, C_1 - C_6 alkyloxycarbonyl, and aryl C_1 - C_6 alkyloxy

carbonyl; R_6 is selected from the group consisting of hydrogen, aryl, and C_1 - C_6 alkyl; with the proviso that when $R_2=R_3=R_5=R_6$ =hydrogen, R_4 is not alkyl.

5

17. A compound of the formula



wherein R_1 and R_3 are selected from the group consisting of hydrogen, hydroxy, C_1 - C_6 alkyloxy, and C_1 - C_6 alkylcarbonyloxy; R_2 is selected from the group consisting of hydrogen, hydroxy, C_1 - C_6 alkyloxy, C_1 - C_6 alkylcarbonyloxy, and C_2 - C_6 alkenyloxy, said alkenyloxy together with the carbon atom of the phenyl ring forming an oxygen heterocycle; and R_4 is selected from the group consisting of phenyl, styryl, phenylbutadienyl, phenylacetylenyl, and $-CH=N$ -phenyl, and substituted phenyl, styryl, phenylacetylenyl, and phenylbutadienyl, wherein the phenyl ring is substituted with 1 to 5 C_1 - C_6 alkyloxy groups; with the provisos that when R_3 is hydrogen, R_1 and R_2 are neither hydroxy nor alkyloxy; when R_1 , R_2 , and R_3 are hydrogen, R_4 is neither phenyl nor alkyloxyphenyl; when R_3 is hydrogen and R_4 is phenyl, neither R_1 nor R_2 is alkylcarbonyloxy; and when R_3 is hydroxy or alkyloxy, R_1 and R_2 are not dihydroxy.

25

18. The compound of claim 17, wherein R_4 is phenyl.

19. The compound of claim 18, wherein R_3 is a C_1 - C_6 alkyloxy.

30

20. The compound of claim 19, wherein R_1 and R_2 are 5,7-di(C_1 - C_6 alkyloxy).

21. The compound of claim 17, wherein R_4 is a 2,4-di(C_1 - C_6 alkyloxy)phenyl.

35

22. The compound of claim 21, wherein R_3 is a C_1-C_3 alkyloxy.

5 23. The compound of claim 22, wherein R_1 and R_2 are the same and are selected from the group consisting of methoxy and ethoxy.

10 24. The compound of claim 17, wherein R_1 is 5-hydroxy and R_2 is one of methoxy and ethoxy.

25. The compound of claim 17, wherein R_4 is phenylacetylenyl.

15 26. The compound of claim 25, wherein R_3 is hydroxy.

27. The compound of claim 26, wherein one of R_1 and R_2 is methoxy.

20 28. The compound of claim 17, wherein said compound is a 4-(C_1-C_6 alkyloxy)-7-styrylvisnagin.

25 29. The compound of claim 28, wherein said compound is selected from the group consisting of 4-methoxy-7-*trans*-styrylvisnagin, 4-ethoxy-7-*trans*-styrylvisnagin, and 4-propoxy-7-*trans*-styrylvisnagin.

30 30. The compound of claim 17, wherein said compound is a C_1-C_6 alkyloxy-7-phenylbutadienylvisnagin.


31. The compound of claim 30, wherein said compound is one of 4-methoxy-7-phenylbutadienylvisnagin and 4-ethoxy-7-phenylbutadienylvisnagin.

35 32. The compound of claim 17, wherein said compound is a C_1-C_6 alkyloxy-7-(CH=N-phenyl)visnagin.

33. The compound of claim 32, wherein said compound is 4-methoxy-7-(CH=N-phenyl)visnagin.

34. The compound of claim 17, wherein said compound is selected from the group consisting of 3,5,7-triacetoxyflavone, 3,5,7-trimethoxyflavone, 3,5,7-triethoxyflavone, 3,7-diethoxy-5-hydroxyflavone, 3,5,7-tripropoxyflavone, 3,4',5,7-tetramethoxyflavone, 2',3,4',7-tetraethoxy-5-hydroxyflavone, 2',3,4',5,7-pentamethoxyflavone, 2',3,4',5,7-pentaethoxyflavone, hexamethylmyricetin, and 3-hydroxy-4'-phenylacetylenyl-6-methoxyflavone.


35. A compound of the formula



wherein R₁ is selected from the group consisting of hydroxy and C₁-C₆ alkyloxy, and M is a divalent radical selected from the group consisting of -CH(OH)-CH(R₂)- and -C(OH)=C(R₂)-, wherein R₂ is selected from the group consisting of styryl and phenylacetylenyl.

36. The compound of claim 35, wherein said compound is selected from the group consisting of 2-phenylacetylenyl-3-hydroxy-6-methoxyflavone, *trans*-2-styryl-3-hydroxy-6-methoxyflavone, and *trans*-2-phenylacetylenyl-3-hydroxy-6-methoxyflavone.


37. A method of treating a mammal comprising selectively blocking one or more adenosine receptors of said mammal by administering to said mammal at least one compound of the formula



- wherein R_1 and R_3 are selected from the group consisting of hydrogen, hydroxy, C_1 - C_6 alkyloxy, and C_1 - C_6 alkylcarbonyloxy;
- 5 R_2 is selected from the group consisting of hydrogen, hydroxy, C_1 - C_6 alkyloxy, C_1 - C_6 alkylcarbonyloxy, and C_2 - C_6 alkenyloxy, said alkenyloxy together with the carbon atom of the phenyl ring forming an oxygen heterocycle; R_4 is selected from the group consisting of phenyl, styryl,
- 10 phenylbutadienyl, phenylacetylenyl, and $-CH=N$ -phenyl, and substituted phenyl, styryl, phenylacetylenyl, and phenylbutadienyl, wherein the phenyl ring is substituted with 1 to 5 C_1 - C_6 alkyloxy groups; with the provisos that when R_3 is hydrogen, R_1 and R_2 are neither hydroxy nor
- 15 alkyloxy; when R_1 , R_2 , and R_3 are hydrogen, R_4 is neither phenyl nor alkyloxyphenyl; and when R_3 is hydrogen and R_4 is phenyl, neither R_1 nor R_2 is alkylcarbonyloxy.

38. A method of treating a mammal comprising
- 20 selectively blocking one or more adenosine receptors of said mammal by administering to said mammal at least one compound selected from the group consisting of genistein, (\pm)dihydrogenistein, sakuranetin, α -naphthoflavone, β -naphthoflavone, amaryllidaceae, oxogalanthine lactam,
- 25 acetylhaemanthine methiodide, 2,3-methylenedioxy-fluorene-9-one, hematoxylin, and arborinine.

39. A compound of the formula



or a pharmaceutically acceptable salt thereof, wherein R₁ is selected from the group consisting of C₁-C₆ alkylcarbonyl, aryl C₁-C₆ alkylcarbonyl, aryl C₂-C₆ alkenylcarbonyl, C₁-C₆ alkyloxy carbonyl, amino C₁-C₆ alkylcarbonyl, and arylcarbonyl, wherein said aryl may be further substituted with halo, nitro, hydroxy, amino or cyano; and R₂ is hydrogen or halogen.

10

40. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a compound of any of claims 1-36 and 39.

15

41. A method of treating a mammal comprising selectively blocking an adenosine receptor of a mammal by administering to said mammal a compound of any of claims 1-36 and 39.

1/17




FIG. 1

2/17




FIG. 2

3/17




FIG. 3

4/17






FIG. 4

5/17

**FIG. 5**

SUBSTITUTE SHEET (RULE 26)

6/17



R₃ = Me, Et, etc. R₄ = alkyl, aryl, etc.

R₅ = Et, MeOEt, Bn, etc. R₆ = H, alkyl, aryl.

FIG. 6

7/17





FIG. 7

8/17

**FIG. 8**

9/17




FIG. 9A

10/17






FIG. 9B


11/17

**FIG. 10A**


12/17

**FIG. 10B**


13/17

**FIG. 11A**


14/17

**FIG. 11B**

15/17

**FIG. 11C**

16/17

**FIG. 11D**

17/17




FIG. 12